



TITLE:

SYNTHESIS OF CELLO-OLIGOSACCHARIDES -  
INFLUENCE OF SUBSTITUENT GROUPS ON  
STEREOSELECTIVE GLYCOSYLATION  
REACTION-( Dissertation\_全文 )

AUTHOR(S):

Takano, Toshiyuki

---

CITATION:

Takano, Toshiyuki. SYNTHESIS OF CELLO-OLIGOSACCHARIDES -INFLUENCE OF  
SUBSTITUENT GROUPS ON STEREOSELECTIVE GLYCOSYLATION REACTION-. 京都大学,  
1990, 農学博士

ISSUE DATE:

1990-03-23

URL:

<https://doi.org/10.14989/doctor.k4576>

RIGHT:

新	制
農	
561	
京大附図	

SYNTHESIS OF CELLO-OLIGOSACCHARIDES  
 —INFLUENCE OF SUBSTITUENT GROUPS  
 ON STEREOSELECTIVE GLYCOSYLATION REACTION—

1 9 9 0

TOSHIYUKI TAKANO

SYNTHESIS OF CELLO-OLIGOSACCHARIDES  
—INFLUENCE OF SUBSTITUENT GROUPS  
ON STEREOSELECTIVE GLYCOSYLATION REACTION—

1 9 9 0

TOSHIYUKI TAKANO

## CONTENTS

LIST OF FIGURES.....	iii
LIST OF TABLES.....	v
LIST OF ABBREVIATIONS.....	vi
INTRODUCTION.....	P1
Chapter 1 Synthesis of starting material for regiospecific glycosylation.....	P6
1-1 Basic synthetic design for cello-oligosaccharide.....	P6
1-2 Synthesis of 2,3,6-tri- <u>Q</u> -substituted glucose derivatives.....	P11
1-2-1 Selection of protective groups.....	P11
1-2-2 Synthesis of acetyl derivatives.....	P12
1-2-3 Synthesis of benzoyl derivatives.....	P22
1-2-4 Glycosylation of acyl derivatives.....	P25
1-3 Summary.....	P26
Chapter 2 Substituent-effect of aglycon on glycosylation reaction.....	P28
2-1 Substituent-effect of acetyl and benzyl groups.....	P28
2-1-1 Synthesis of 3- <u>Q</u> -acetyl derivatives.....	P29
2-1-2 Synthesis of 3- <u>Q</u> -benzyl derivatives.....	P31
2-1-3 Glycosylation of 3- <u>Q</u> -acetyl and 3- <u>Q</u> -benzyl derivatives.....	P35
2-2 Substituent-effect of other alkyl and acyl groups.....	P38
2-2-1 Substituent-effect of 3- <u>Q</u> -alkyl group.....	P39
2-2-2 Substituent-effect of 3- <u>Q</u> -acyl group.....	P43
2-3 Summary.....	P47
Chapter 3 Substituent-effect of glycon on glycosylation	

reaction.....	P50
3-1 Substituent-effect of $\alpha$ -imide on glycosylation	
reaction.....	P50
3-1-1 Synthesis of $\alpha$ -imides.....	P52
3-1-2 Stability of $\alpha$ -imides and their	
glycosylation.....	P54
3-2 Conclusive selection of starting material for	
cello-oligosaccharide synthesis.....	P58
3-3 Summary.....	P64
CONCLUSIONS.....	P66
EXPERIMENTAL.....	P69
REFERENCES.....	P114
ACKNOWLEDGEMENTS.....	P127

## LIST OF FIGURES

Figure 1.	Linear synthetic method for cello-oligosaccharides.....	P8
Figure 2.	Convergent synthetic method for cello-oligosaccharides.....	P9
Figure 3.	Synthetic route for compound 9 from compound 10.....	P13
Figure 4.	Reaction mechanism for tritylation with trityl chloride / pyridine.....	P15
Figure 5.	Reaction mechanism of the formation of compound 24 from compound 17.....	P20
Figure 6.	Synthetic route for compound 30 from compound 14 or 28.....	P23
Figure 7.	Synthetic route for compounds 17 and 39 from compound 11.....	P30
Figure 8.	Synthetic route for compounds 40 and 41 from compound 44.....	P33
Figure 9.	Synthetic route for compounds 55-57 from compound 58.....	P40
Figure 10.	Synthetic route for compounds 66-69 from compound 13.....	P45
Figure 11.	Synthetic route for compounds 83, 84, 88 and 90 from compound 25.....	P53
Figure 12.	Synthetic route for compounds 85-87 and 89 from compounds 40, 106 and 113.....	P55
Figure 13.	Synthetic route for compounds 122 and 123 from compound 44.....	P61

Figure 14. Synthetic route for compound 128

from compound 122.....P63

## LIST OF TABLES

Table 1.	Results of glycosylation reactions between $\alpha$ -imide 34 and aglycons 17, 39-41.....	P36
Table 2.	Results of glycosylation reactions between $\alpha$ -imide 34 and aglycons 40, 55-57.....	P42
Table 3.	Results of glycosylation reactions between $\alpha$ -imide 34 and aglycons 17, 66-69.....	P46
Table 4.	Results of the stability and reactivity of the $\alpha$ -imides 34, 83-90.....	P56



## LIST OF ABBREVIATIONS

Ac	= acetyl	$-\text{COCH}_3$
All	= allyl	$-\text{CH}_2\text{CH}=\text{CH}_2$
Bz	= benzoyl	$-\text{COC}_6\text{H}_5$
Bzl	= benzyl	$-\text{CH}_2\text{C}_6\text{H}_5$
Car	= carbamoyl	$-\text{CONHC}_6\text{H}_5$
DBU	= 1,8-diazabicyclo[5.4.0]-7-undecene	
DMF	= N,N-dimethylformamide	
DMSO	= dimethyl sulfoxide	
D.P.	= degree of polymerization	
DPM	= diphenylmethyl	$-\text{CH}(\text{C}_6\text{H}_5)_2$
MCA	= monochloroacetyl	$-\text{COCH}_2\text{Cl}$
eq	= equivalent	
MOM	= methoxymethyl	$-\text{CH}_2\text{OCH}_3$
MTM	= methylthiomethyl	$-\text{CH}_2\text{SCH}_3$
m.p.	= melting point	
Ph	= phenyl	$-\text{C}_6\text{H}_5$
Piv	= pivaloyl	$-\text{COC}(\text{CH}_3)_3$
pMBzl	= p-methoxybenzyl	$-\text{CH}_2\text{C}_6\text{H}_4\text{OCH}_3$
P-TLC	= preparative TLC	
p-TsOH	= p-toluenesulfonic acid	
Rf	= distance travelled by zone, divided by distance travelled by solvent front	
r.t.	= room temperature	
TMSCl	= trimethylsilyl chloride	
Tr	= trityl	$-\text{C}(\text{C}_6\text{H}_5)_3$
triflate	= trifluoromethanesulfonate	

## INTRODUCTION

Cello-oligosaccharides are generally defined as oligosaccharides consisting of two to ten D-glucopyranosyl residues jointed through  $\beta$ -1,4-glucosidic bonds which are readily hydrolyzed under acidic conditions to yield the constituent glucose.<sup>76)</sup> These cello-oligosaccharides are not present free in nature but present only as structural units of cellulose.

Since Payen named "cellulose" for a polymer of D-glucose as a cell wall component of the higher plants in 1838, many studies on cellulose have been conducted for 150 years up to date.<sup>60)</sup> Nevertheless, there still remains several fundamental problems related to cellulose chemistry such as mechanisms of biosynthesis and biodegradation, crystal structure, mechanism of mercerization, relationship between chemical structure of cellulose derivatives and their physicochemical properties and so on. These problems may be clearly solved by the use of cellulose model compounds such as cello-oligosaccharides. Biodegradation mechanism for lignin as a complex macromolecule recently, has been elucidated by the use of low molecular weight model compounds "oligo-lignols".<sup>35)</sup> Hence, cello-oligosaccharides may play an important role in the future on fundamental cellulose research.<sup>64)</sup>

These cello-oligosaccharides are obtainable by both partial degradation of cellulose and chemical synthesis from D-glucose.<sup>106)</sup> Since cellobiose octaacetate was first

prepared by Franchimont<sup>17)</sup> in 1879, many researchers reported the preparation of the cello-oligosaccharides from cellulose by acetolysis, acid-catalyzed hydrolysis and enzymatic hydrolysis.<sup>106)</sup> Miller et al.<sup>63)</sup> prepared a series of cello-oligosaccharides up to celloheptaose in 1960. However, the yield of the cello-oligosaccharides by the partial degradation was extremely low for oligomers with high molecular weight, and isolation and purification of a series of the cello-oligosaccharide are tedious and not so easy.

On the other hand, chemical synthesis could be reliable and useful to obtain a pure cello-oligosaccharide, although there are many synthetic problems such as selections of the protective groups and glycosylation reaction to be solved.

The first synthesis of cellobiose was carried out by Freudenberg and Nagai<sup>19)</sup> in 1933 and thereafter many studies have been reported on the synthesis of cellobiose and cellotriose.<sup>31),95)</sup> Cellotetraose was synthesized by Schmidt and Michel<sup>82)</sup> in 1982 and by Takeo et al.<sup>96)</sup> in 1983. However, higher cello-oligosaccharides than tetramer have not yet been synthesized and will not be obtained readily by their methods because these methods require many reaction steps. At present, the general synthetic method for cello-oligosaccharides has not been established.

The first synthetic approach toward cellulose was tried by Schlubach and Luhrs<sup>81)</sup> in 1941. Since then, many researchers have attempted to synthesize cellulose. In spite

of the enormous effort devoted to cellulose synthesis, there has been no paper which completely solved the synthetic problems for cellulose.<sup>70),71),94)</sup> Therefore, the synthetic methods tried so far for cellulose can not be directly applied for the synthesis of cello-oligosaccharides, although some useful information could be obtained.

In principle, there are three problems to be solved for the synthesis of a cello-oligosaccharide; (1) regiospecific control: glycosylation of the hydroxyl group at the O-4 position of the glucopyranosyl residue has to be performed, (2) stereospecific control: glucosidic linkage of glucopyranosyl residues of  $\beta$ -configuration has to be made, (3) control of degree of polymerization (D.P.): the desired sequence of units has to be formed. The first problem could be solved by use of 2,3,6-tri-O-substituted glucopyranose derivative as a starting material, the second by the stereospecific glycosylation reaction, and the third by the stepwise condensation method. However, the main obstacle in the synthesis lies in the second problem, glycosylation reaction.<sup>73),107)</sup>

Newly-developed glycosylation methods and extensive data on glycosylation reactions have made possible to prepare several complex oligosaccharides such as hexa- and nona-saccharides fragments of the O-antigenic polysaccharide,<sup>15)</sup> heparin-like penta-saccharide,<sup>51)</sup> xylo- and malto-oligosaccharides.<sup>49),80)</sup> However, the glycosylation reaction is complicated and there still remain many problems for the highly stereoselective glycosylation with high yield, such

as selection of the leaving groups at anomeric center, activating reagents, solvents used and so on. Although there are no systematic studies from this point of view, the effects of the protective groups of both aglycon (hydroxyl component) and glycon (glycosyl component) on the stereoselective glycosylation reactions have been considered extremely important.

Substituent effect of the aglycon on glycosylation of N-acetyl glucosamine derivative by Koenigs-Knorr method has been reported by Sinay.<sup>87)</sup> However, it is not known whether the result of Sinay<sup>87)</sup> is applicable to cello-oligosaccharides synthesis.

Paulsen<sup>73),74)</sup> discussed the influence of glycon protective groups in the synthesis of  $\beta$ -D-manno- and  $\beta$ -L-rhamnoglycosidic derivative. While Boeckel et al.,<sup>3),4),5)</sup> reported substituent-effects of  $\alpha$ -D-gluco- and  $\alpha$ -D-manno-pyranosyl bromides on glycosylation by insoluble salts. However, the results obtained by Paulsen is not necessarily agreed with those by Boeckel et al.. It is uncertain whether these results are applicable to glycosylation method other than Koenigs-Knorr method.

Therefore, a systematic investigation of the effect of the protective groups of aglycon and glycon on  $\beta$ -glycosylation is needed in the cello-oligosaccharides synthesis.

In the present thesis, the author focused on the influence of substituent groups of the aglycon and glycon on

glycosylation reaction (especially,  $\beta$ -1,4 glucosidic bond formation) for the synthesis of cello-oligosaccharides.

Chapter 1 contains a general synthetic design of cello-oligosaccharides and the synthesis of acyl 2,3,6-tri-O-substituted glucose derivative and its glycosylation according to this design.<sup>67),89),90)</sup>

In Chapter 2, the synthesis of aglycons with various protective groups and the influence of the substituent groups of the aglycon on glycosylation reaction are described.<sup>90),92)</sup>

In Chapter 3, the synthesis of glycons with different substituent pattern and the substituent effect of the glycon on glycosylation reactions are described.<sup>91)</sup> In addition, the selection of the most suitable starting material for the synthesis of cello-oligosaccharides and the future possible usage of the compound is discussed.<sup>93)</sup>

The detailed experimental methods and data of compounds are summarized in EXPERIMENTAL.

## Chapter 1 Synthesis of starting material for regiospecific glycosylation

### 1-1. Basic synthetic design for cello-oligosaccharide

Nakatsubo et al.<sup>67)</sup> proposed a synthetic design of cello-oligosaccharides from D-glucose as shown in Figs. 1 and 2.

In the synthetic design the substituted glucose derivative 1, which has three kinds of protective groups, X-, Y- and R-groups, is required as a starting material for the regulation of regiospecificity. Here, X- and Y-groups are "temporary" protective groups, and R-group is a "persistent" protective group.<sup>18),27)</sup> The temporary protective groups should be removed independently without giving any influence on the other functional groups before glycosylation reaction. It is prerequisite for the selection of these three protective groups that the Y- and R-groups or X- and R-groups must not be changed on the removal of the temporary protective X- or Y-groups, respectively. The persistent protective group is removed at the final step of the synthesis for the cello-oligosaccharide with the expected degree of polymerization (D.P.). Therefore, the R-group should be fairly stable under various conditions, but the removal conditions do not cause decomposition of the hydroxyl groups or  $\beta$ -glucosidic bonds in the synthesized cello-oligosaccharides. It should be considered that the selection of the R-group determines the choice of X- and Y-groups and vice versa.

The synthesis of cello-oligosaccharides is conducted from the starting material 1 by two reactions; deprotection of X- or Y-groups and  $\beta$ -glycosylation. The removal of X- or Y-groups from the starting material 1 gives glycon 2 or aglycon 3, respectively.  $\beta$ -Glycosylation between glycon 2 and aglycon 3 leads to the formation of cellobiose derivative 4 which has the same protective system as starting material 1. Cellobiose derivative 4 is further converted into glycon 5 and aglycon 6 by deprotection of X- or Y-groups, respectively. Here, two major synthetic methods are conceivable for the synthesis of cello-oligosaccharides; a linear synthetic method and a convergent synthetic method.<sup>102),103)</sup> In the former method (Fig. 1),  $\beta$ -glycosylation is performed between glycon 5 and aglycon 3 or between glycon 2 and aglycon 6 to afford cellotriose derivative 7 which also has the same protective system as compound 1. Compound 7 is combined with compound 2 or 3 to give cellotetraose derivative 8. After repeating the above two reactions, a series of cello-oligosaccharide (whose D.P.= 1,2,3,4....) could be theoretically obtained in this method. In the latter method (Fig. 2),  $\beta$ -glycosylation is conducted between glycon 5 and aglycon 6 to give cellotetraose derivative 8 with the same protective system as compound 1. Further, the two compounds 8 are combined to afford the cellooctaose derivative. Thus, in this method, a series of cello-oligosaccharide (whose D.P.= 1,2,4,8....) could be prepared theoretically by repeating of the two






Figure 1. Linear synthetic method for cello-oligosaccharides.




Figure 2. Convergent synthetic method for cello-oligosaccharides.

reactions.

Since the first synthesis of an O-glycoside was carried out by Michael in 1879,<sup>62)</sup> a number of methods for the synthesis of O-glycoside have been proposed,<sup>2),40),48),101)</sup> i.e., alcoholysis of free sugars (Fischer method) and syntheses from 1-halo sugars (Koenigs-Knorr method), from 1-O-acyl sugars (Helferich method), from 1,2-orthoesters (Orthoester method), from 1-fluoro sugars (Fluoride method), from 1-imidate sugars (Imidate method) and so on. However, no universal glycosylation methods have been established for the oligosaccharide synthesis so far.

The trichloroacetimidate method (hereafter imidate method) was selected in the present investigation for the following four reasons;<sup>83),84)</sup> [1] Glycosylation can be conducted under a very mild reaction condition (for example, at  $-78^{\circ}\text{C}$  for about 30 min). [2] A  $\text{S}_{\text{N}}-2$  reaction mechanism has been proposed to give only the  $\beta$ -glycoside from the  $\alpha$ -imidate at low temperature with boron trifluoride etherate catalysis. [3] By the selection of catalysts and protective groups, reactivities that are comparable to those obtained for the glycosyl halide / silver triflate system (Koenigs-Knorr method) can be expected. [4] Schmidt and Michel<sup>82)</sup> used this method for the synthesis of cellotetraose derivative and Kawada et al.<sup>44)</sup> successfully synthesized the cellobiose derivative in high yield.

In the present synthetic design, synthesis of the substituted glucose derivative 1 (namely, the introduction and removal of the protective groups) and  $\beta$ -glycosylation

reaction are very important.

## 1-2. Synthesis of acyl 2,3,6-tri-O-substituted glucose derivatives

### 1-2-1. Selection of protective groups


In the previous section, the basic synthetic design for cello-oligosaccharides was proposed. The first step in this design is the synthesis of the substituted glucose derivative 1.

Acetyl group was selected as the persistent protective R-group for the following advantages in this investigation. They are; [1] It could be easily introduced and removed. [2] The acetyl group introduced into the O-2 position of a glucose derivative (glycon) is expected to have a favorable effect on the formation of  $\beta$ -glucosidic bond by way of neighbouring group participation.<sup>2)</sup> [3] Acetyl glucose derivatives have good solubility in organic solvents and could be easily crystallized. [4] Acetyl glucose derivatives are relatively easy to determine their chemical structure by their NMR spectra analysis. [5] Enough information on cellulose acetate is available, because it is widely used in the field of cellulose chemistry. However, the acetyl group also has several drawbacks such as the possible intramolecular migration, the instability under basic conditions and the lack of UV absorption.

On the other hand, 2,2,2-trichloroethyl and benzyl groups were chosen as the X- and Y-groups, respectively.


2,2,2-Trichloroethyl group of the sugars with acetyl and benzyl substituents can be selectively removed by zinc powder in acetic acid, and was used for the protection of the anomeric hydroxyl group by Lemieux and Driguez.<sup>55),56)</sup> Benzyl group can be deblocked by catalytic hydrogenolysis independently. Benzyl derivative could be observed on TLC under UV-light. This makes it possible to easily follow progress of the reaction and to purify the expected compound by TLC. Kovac<sup>49)</sup> used benzyl group for the protection of C-4 hydroxyl group in the synthesis of xylo-oligosaccharides.

Thus, 2,2,2-trichloroethyl 2,3,6-tri-O-acetyl-4-O-benzyl- $\beta$ -D-glucopyranoside (**9**) was selected as a starting material 1.



### 1-2-2 Synthesis of the acetyl derivatives

The synthetic route for compound **9** from D-glucose (**10**) is shown in Fig. 3. The key compound of this route is compound **17** which has a free hydroxyl group at the C-4 position of D-glucose. Many synthetic methods for such glucose derivatives, e.g. acyl group migration,<sup>6),33)</sup> selective acylation,<sup>86),100)</sup> reductive cleavage of 4,6-O-benzylidene derivative<sup>24),25)</sup> and use of a 4-O-nitro



- a)  $\text{Ac}_2\text{O}$  /  $\text{AcONa}$  /  $80^\circ\text{C}$  / 17hr    b)  $\text{CCl}_3\text{CH}_2\text{OH}$  /  $\text{BF}_3\text{-Et}_2\text{O}$  /  $\text{CH}_2\text{Cl}_2$  /  $0^\circ\text{C}$  / 15hr  
 c) 28%- $\text{NaOMe}$  in  $\text{MeOH}$  /  $\text{MeOH}$  /  $0^\circ\text{C}$  / 1hr    d)  $\text{TrCl}$  /  $\text{Et}_3\text{N}$  / anhydrous dioxane /  $100^\circ\text{C}$  / 1hr  
 e)  $\text{Ac}_2\text{O}$  /  $\text{AcONa}$  /  $100^\circ\text{C}$  / 1hr    f) and g)  $p\text{-TsOH}$  /  $\text{CH}_2\text{Cl}_2$  / r.t. / 3hr  
 h)  $\text{BzIOC(=NH)CCl}_3$  /  $\text{CF}_3\text{SO}_3\text{H}$  / anhydrous  $\text{CH}_2\text{Cl}_2$  / anhydrous cyclohexane / r.t. / 5.5hr

Figure 3. Synthetic route for compound 9 from compound 10.

intermediate,<sup>32)</sup> have been reported. Considering their yields, the number of reaction steps and simplicity of the preparation, a modified acyl group migration was applied to obtain compound 17 in this investigation.

Compound 12 was prepared from glucosepentaacetate (11) which was a mixture of  $\alpha$ - and  $\beta$ -anomers (about 1:5) prepared by acetylation of compound 10 with acetic anhydrous / sodium acetate. First, Koenigs-Knorr reaction between tetra-O-acetyl- $\alpha$ -D-glucopyranosyl bromide, which was synthesized from compound 11 by the treatment with 25% hydrogen bromide acetic acid solution, and 2,2,2-trichloroethanol was examined in the presence of a silver or mercury salt such as silver carbonate, silver sulfate and mercury (II) bromide. However, yield of the expected glycoside was about 20%, probably because of poor nucleophilicity resulting from the large electron-attracting effects of the three chlorine atoms and large steric hindrance of 2,2,2-trichloroethanol. Then, a modified Helferich method<sup>65)</sup> was tried. Compound 11 and 2,2,2-trichloroethanol were fused in the presence of catalytic amounts of p-toluenesulfonic acid at 100°C under 100mmHg to afford the expected glycoside in about 40% yield. Finally, the method reported by Magunusson et al.<sup>59)</sup> was found to be the best method in respect to the yield and simplicity of the reaction. The solution of peracetyl glucose 11 and 2,2,2-trichloroethanol was treated with boron trifluoride etherate at room temperature for about 12 hours. After recrystallization, compound 12 was obtained in about 60% yield. This method was also useful for large scale

experiments using 78g (0.2 mol) of the starting material 11.

Compound 13, obtained by the deacetylation of compound 12 with sodium methoxide, was tritylated by the use of tritylchloride / pyridine system in a standard manner<sup>34)</sup> to afford trityl ether 14 in a maximum of 67% yield, but the reproducibility of this method was poor. A possible reaction mechanism for the tritylation by this method is shown in Fig. 4.

Chaudhary and Hernandes<sup>8)</sup> suggested that the formation of the N-tritylpyridinium salt (18) is the rate determining step in this reaction. Furthermore, it was found that the treatment of compound 14 with pyridinium hydrochloride (19)




Figure 4. Reaction mechanism for tritylation with trityl chloride / pyridine.




in pyridine gave the starting compound 13. This experimental result indicates that this reaction is reversible as shown in Fig. 4.

Therefore, the following three points are extremely important to increase the reaction rate and to shift the equilibrium toward the product: [1] The use of a base stronger than pyridine (e.g. 4-dimethylaminopyridine, triethylamine, etc.) is recommended for accelerating the formation of the salt 18. [2] The use of a solvent in which compound 19 is insoluble will cause compound 19 to precipitate and will force the reaction to completion. [3] Anhydrous conditions are essential to avoid the hydrolysis of tritylchloride and the expected compound 14. Keeping in mind the above points, the synthesis of compound 14 was attempted using many combinations of bases and solvents. Among these, the triethylamine / dioxane system was found to be the best. When compound 13 was heated with triethylamine in anhydrous dioxane at 100°C for 1 hour, triethylamine hydrochloride was effectively removed from the reaction system as a precipitate, and then compound 14 was obtained in 86.9% yield. This method was found to be the most satisfactory method, with respect to both reproducibly high yield and simplicity of the preparations. The purification of trityl ether 14 by silica gel column chromatography should be carried out as rapidly as possible, because silica gel catalyzed detritylation has been reported.<sup>54)</sup>

In general, trityl ethers are cleaved by a number of methods, including acidic hydrolysis and catalytic,

chemical, or electrolytic reduction.<sup>28)</sup> Here, acid-catalyzed reactions using a Lewis acid or a protonic acid were examined. The use of boron trifluoride etherate gave preferentially the detritylation compound 16 with the production of only a small amount (less than 5% yield) of compound 17 resulted from acetyl group migration. On the other hand, when compound 15 was treated with p-toluenesulfonic acid, both the detritylation and subsequent acetyl group migration proceeded smoothly; compound 17 was obtained in 86.8% yield without any residual amount of compound 16. Similar results were obtained using Amberlyst H-15 (H<sup>+</sup>) or DL-camphor-10-sulfonic acid, but the reaction proceeded very slowly. Thus, the detritylation with a Lewis acid gave preferentially the detritylation compound 16, whereas the detritylation with a protonic acid afforded exclusively compound 17.


These results could be attributed to the coordination of boron trifluoride with the ring oxygen and O-6 as shown in intermediate 20. Such coordination would reduce the electron density on O-6 and inhibit the free rotation of the C-6



hydroxyl group around the C<sub>5</sub>-C<sub>6</sub> linkage, thus preventing the formation of the desired cyclic orthoester intermediate 21, from which compound 17 was derived. It seems that this coordination is similar to that of zinc bromide in detritylation<sup>47),61)</sup> or demethoxymethylation.<sup>10)</sup>

All of the above detritylations were carried out in methylene chloride. In 20% methanol-methylene chloride with the presence of a Lewis acid or a protonic acid, acetyl group migration did not occur at all. The detritylation compound 16 was obtained in over 95% yield in both cases. This result is interesting, but a satisfactory explanation is not available at this time.

When compound 17 was treated with benzyl bromide / silver oxide in N,N-dimethylformamide (DMF), which is a typical benzylation method in carbohydrate chemistry,<sup>28)</sup> the expected benzyl ether 9 was obtained in only 20% yield with several by-products. The low yield is attributed to the low reactivity of the C-4 hydroxyl group,<sup>46)</sup> and to the possibility of acetyl group migration under these reaction conditions.<sup>29),88)</sup> An alternative method which involves acetyl group migration from O-4 to O-6 and simultaneous benzylation<sup>85)</sup> was applied to compound 16, but the expected compound 9 was obtained in only 20% yield. The benzylation of compound 17 with benzyl bromide / sodium hydride at 0°C for a short time (30 min) gave compound 9 in 17% yield with many by-products. Interestingly, 2,2,2-trichloroethyl 3,4,6-tri-O-acetyl-2-O-benzyl-β-D-glucopyranoside (24) and 2,2,2-trichloroethyl 2,4,6-tri-O-acetyl-3-O-benzyl-β-D-



glucopyranoside (25) were identified as by-products. The conversion of the compound 17 to the compound 24 indicates, that intramolecular acetyl group migration from the O-2 position to the O-4 position via an intermediate 22 taking 1C-conformation occurs and is then followed by the benzylation of the resulting free hydroxyl group at the C-2 position of the intermediate 23 as shown in Fig. 5.

All of the above investigations suggest that the benzylation of the C-4 hydroxyl group cannot be carried out in high yield under neutral or basic conditions, and that the introduction of a protective group is only possible under acidic conditions. Then, the benzylation by a Friedel-Crafts type reaction<sup>79)</sup> and treatment with the benzyltrichloroacetimidate / trifluoromethanesulfonic acid system (benzylimidate method)<sup>41), 104)</sup> were examined. Compound 17 was treated with benzyl alcohol / phosphorus pentaoxide ( $P_2O_5$ ), but the expected benzyl ether 9 was only obtained in a maximum of 20% yield, and the results were not reproducible. On the other hand, compound 17 was benzylated by the benzylimidate method, and the yield of compound 9 increased to about 50%. At present, no benzylation method





Figure 5. Reaction mechanism of the formation of compound **24** from compound **17**

has been found for compound 17 superior to the benzylimidate method.

Since the introduction of the benzyl group into the C-4 hydroxyl group was so difficult, methylthiomethyl (MTM),<sup>77),78)</sup> diphenylmethyl (DPM),<sup>52)</sup> or methoxymethyl (MOM)<sup>20)</sup> group was examined as the Y-group. The MTM, DPM, and MOM groups could be introduced by Friedel-Crafts type reactions and removed selectively by the use of mercury chloride, neutral hydrogenolysis, or a catalytic amount of hydrochloric acid, respectively.

Methylthiomethylation of compound 17 with acetic anhydride / acetic acid / dimethylsulfoxide (DMSO) gave the MTM ether 26 in 63.4% yield. But the MTM group was unstable in the presence of zinc powder / acetic acid, with which the trichloroethyl group was removed,<sup>28)</sup> compound 26 was not suitable for the purpose. However, the MTM group is a useful protective group for the C-4 hydroxyl group of glucose.

Treatment of compound 17 with benzhydrol / P<sub>2</sub>O<sub>5</sub> or methyral / P<sub>2</sub>O<sub>5</sub> gave the DPM ether 27 or the MOM ether 28 in 70.3% or 87.6% yield, respectively. The satisfactory yield from these reactions indicates that the respective



26 Y = -CH<sub>2</sub>SCH<sub>3</sub> (-methylthiomethyl, -MTM)

27 Y = -CH(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub> (-diphenylmethyl, -DPM)

28 Y = -CH<sub>2</sub>OCH<sub>3</sub> (-methoxymethyl, -MOM)

cations may be produced readily and that side reactions do not occur appreciably under these reaction conditions. Thus, these results indicate that the introduction of a functional group into C-4 hydroxyl group of an acetylated glucose derivative proceeds in high yield only under acidic conditions.

### 1-2-3 Synthesis of the benzoyl derivatives

Either compound 27 or 28 described above is thought to be useful as a starting material 1 for the synthesis of cello-oilgosaccharides. However, the absence of UV absorption in compound 28 is inconvenient to follow the progress of a reaction and to purify the expected compound on TLC. Therefore, the introduction of benzoyl group instead of acetyl group as the R-group was examined. The merits of the benzoyl group are similar to those of the acetyl group, although the former has greater steric hindrance than the latter. The synthesis of 2,2,2-trichloroethyl 2,3,6-tri-O-benzoyl-4-O-methoxymethyl- $\beta$ -D-glucopyranoside (30) was carried out via two synthetic routes as shown in Fig. 6.

First, compound 30 was synthesized from compound 28. Deacetylation of compound 28 with sodium methoxide, followed by benzoylation with benzoyl chloride / anhydrous pyridine gave compound 30 in 63.5% yield.

Then, compound 30 was prepared from compound 14 by the method similar to that used for the synthesis of compound 28. Compound 14 was converted into compound 31 in 87.1%





Figure 6. Synthetic route for compound **30** from compound **14** or **28**.



yield by the use of benzoyl chloride / anhydrous pyridine. In general the benzoyl group migrates less readily than the acetyl group.<sup>1)</sup> Indeed, the detritylation of compound 31 with p-toluenesulfonic acid in dichloromethane at room temperature gave 2,2,2-trichloroethyl 2,3,4-tri-O-benzoyl-β-D-glucopyranoside in 63.7% yield. Benzoyl group migration did not occur. However, when compound 31 was treated with p-toluenesulfonic acid in 1,2-dichloroethane at 50°C, the benzoyl group migrated from O-4 to O-6 to give compound 32 in 70.7% yield. Thus, the benzoyl group is better than the acetyl group as the R-group for the synthesis of a glucose derivative which has a free hydroxyl group at the C-6 position. The methoxymethylation of compound 32 with methyral / P<sub>2</sub>O<sub>5</sub> gave compound 30 in 77.5% yield.

Finally, the introduction of the monochloroacetyl group into the C-4 hydroxyl group was conducted. The monochloroacetyl (MCA) group is useful because it is selectively removed by thiourea but is resistant to acidic conditions which are necessary for the bromination of the anomeric hydroxyl group in Koenigs-Knorr type glycosylation.<sup>9),28)</sup> Compound 32 was converted into the MCA ether 33 in 81.0% yield by the use of chloroacetic anhydride / pyridine.<sup>23)</sup>




#### 1-2-4 Glycosylation of acyl derivatives

The second step in the design described in Section 1-1 is the synthesis of cellobiose derivative by the use of the synthesized glucose derivative.

Kawada et al.<sup>44)</sup> reported that the glycosylation between 2,3,6-tri-O-benzyl-4-O-p-methoxybenzyl- $\alpha$ -D-glucopyranosyl trichloroacetimidate (**34**) and allyl 2,3,6-tri-O-benzyl- $\beta$ -D-glucopyranoside (**35**) gave the corresponding cellobiose derivative **36** in 83.7% yield. Then, glycosylation of aglycon **17** with glycon **34** was made under the same reaction conditions reported by Kawada et al.<sup>44)</sup>

Imidate **34** and aglycon **17** were dissolved in anhydrous dichloromethane containing molecular sieves 4A; then the solution was cooled below  $-70^{\circ}\text{C}$ ; boron trifluoride etherate in anhydrous dichloromethane was added; after 2 hours, the reaction mixture was worked-up and purified by TLC to give



compound 37 and compound 38 in 16.4% and 29.0% yield, respectively.

The products were identified by  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR spectroscopies. The characteristic signals, derived from the 2,2,2-trichloroethyl and p-methoxybenzyl groups in the  $^1\text{H}$ -NMR spectra of compounds 37 and 38, provide the structural proof that glycosylation occurred between glycon 34 and aglycon 17. The peaks of compound 37 that appear at 4.26 (1H, d,  $J=8.0$ ) and 102.8 (C-1') conclusively indicate the  $\beta$ -configuration. And the peaks of compound 38 that appear at 4.80 (1H, d,  $J=3.5$ ) and 99.1 (C-1') also indicate the  $\alpha$ -configuration.

This result obviously indicates that the glycosylation is tremendously affected by the protective groups pattern of the aglycon, and that aglycon 17 is unsuitable in the glycosylation by the imidate method. Therefore, before the synthesis of cello-oligosaccharides, it is required to examine the effect of the protective groups of the aglycon and glycon on glycosylation reaction in order to select the best protective group pattern.

### 1-3 Summary

For the synthesis of cello-oligosaccharide, regiospecific and stereospecific glycosylation reaction should be conducted. In this chapter, the basic prerequisite of starting material was discussed at first. The substituted glucose derivative should be protected by three kinds of protective groups, X-, Y- and R-groups as a starting

material (Section 1-1).

Then, 2,2,2-trichloroethyl 2,3,6-tri-O-acyl- $\beta$ -D-glucopyranoside derivatives, which were satisfied with the basic prerequisite, were selected as a starting material and their synthetic route was investigated (Section 1-2-1 to 1-2-3). In the course of these syntheses, it turned out that trityl chloride / triethylamine / dioxane system was the most satisfactory method for tritylation with respect to the yield and the simplicity of preparation (Fig. 3, step d). It was also found that acetyl group migration occurred with a protonic acid, but did not with a Lewis acid (Fig. 3, step g). These results will give useful information in cellulose chemistry, because the tritylation and detritylation reactions are very important reaction for the preparation of regioselective substituted cellulose derivatives, and acetyl derivative is quite common in cellulose chemistry.


Next, the synthesis of cellobiose derivative was tried by the use of the acyl derivative synthesized (Section 1-2-4). However, contrary to the result reported by Kawada et al., the glycosylation of acyl glucose derivative by the imidate method gave a mixture of  $\beta$ - and  $\alpha$ -glycosides. This result obviously indicated that the stereochemistry of the product on glycosylation reaction was tremendously affected by the protective group pattern of aglycon.

## Chapter 2 Substituent-effect of aglycon on glycosylation reaction

### 2-1 Substituent-effect of acetyl and benzyl groups

In Chapter 1 (Section 1-2-4), it was described that the glycosylation is greatly influenced by the substituent pattern of protective groups in aglycon. Sinay<sup>87)</sup> reported that the presence of C-3 ether substituent of the aglycon was very effective for increasing the yield of  $\beta$ -D-(1  $\rightarrow$  4) linked disaccharide derivatives. Takeo *et al.*<sup>96)</sup>, in consideration of the results obtained by Sinay<sup>87)</sup>, recently reported the high yield syntheses of cellobiose, cellotriose and cellotetraose. On the other hand, Paulsen<sup>73),74)</sup> reported that alkyl substituted derivative are more reactive than those bearing acyl substituents. But, no systematic investigation has been reported for the substituent-effect of aglycon on stereospecific glycosylation reaction.

In this section, four aglycons 17, 39, 40 and 41 were selected for the glycosylation reaction with glycon 34. The C-3 and C-6 hydroxyl groups of these aglycons were protected with different combinations of the electron-donating benzyl



group and the electron-withdrawing acetyl group in order to investigate the effect of these protective groups on the anomer ratio in the glycosylation reactions with glycon 34.

### 2-1-1 Synthesis of 3-O-acetyl derivatives

The synthetic route for aglycons 17 and 39 from glucopentaacetate (11) is shown in Fig.7. The synthesis of compound 17 was described in detail in the previous chapter (Chapter 1, Section 1-2-2). The key reaction in the synthesis of compound 39 was the reductive cleavage of 4,6-O-benzylidene glucopyranoside (Fig. 7, step h).

Garegg et al.<sup>24),25)</sup> reported that the highly regioselective, reductive cleavage of 4,6-O-benzylidene hexopyranoside with sodium cyanoborohydride ( $\text{NaCNBH}_3$ ) -  $\text{HCl}(\text{gas})$  in tetrahydrofuran gave the 6-O-benzyl ether, which has a free hydroxyl group at the C-4 position, in high yield. On the other hand, Johansson and Samuelsson<sup>42)</sup> reported that the reductive ring-opening of 4,6-O-p-methoxybenzylidene hexopyranosides with  $\text{NaCNBH}_3$ -trifluoroacetic acid ( $\text{CF}_3\text{COOH}$ ) in  $N,N$ -dimethylformamide (DMF) or with  $\text{NaCNBH}_3$ -trimethylsilyl chloride ( $\text{TMSCl}$ ) in acetonitrile gave the 6- or 4-O-p-methoxybenzyl ether derivative, respectively, in high yield with good regioselectivity.

Since the use of  $\text{HCl}(\text{gas})$  requires tedious handling, the  $\text{NaCNBH}_3$ - $\text{CF}_3\text{COOH}$  and  $\text{NaCNBH}_3$ - $\text{TMSCl}$  systems were examined in the present investigation. Compound 43 was treated with  $\text{NaCNBH}_3$ - $\text{CF}_3\text{COOH}$  in DMF for 12 hours to afford only compound




Figure 7. Synthetic route for **17** and **39** from compound **11**.

39 in 60% yield, as expected. However, different from the results of Johansson and Samuelsson<sup>42)</sup> treatment with  $\text{NaCNBH}_3$ - $\text{TMSCl}$  in acetonitrile for 3 hours also gave only compound 39 in 60% yield. That is, the attack of either electrophile ( $\text{CF}_3\text{COOH}$  or  $\text{TMSCl}$ ) occurred only on the 0-4 position of the benzylidene group to result in the formation of only 6-0-benzyl ether 39 without any steric effect of the electrophile.

It has been reported that the regioselectivity from the reductive ring-opening of a benzylidene acetal is associated with the steric hindrance exerted by the C-3 substituent<sup>57)</sup>. However, compound 49 with a bulky benzyl group at the 0-3 position also gave only 6-0-benzyl ether 41 by the  $\text{NaCNBH}_3$ - $\text{TMSCl}$  treatment, as described in the latter section (Fig. 8). Thus, the regioselectivity seems to also depend on the nature of the benzylidene used. For p-methoxybenzylidene kinetic control favors the attack of  $\text{TMSCl}$  on the 0-6 position, resulting in the production of the 4-0-p-methoxybenzyl ether, but for benzylidene thermodynamic control favors the attack of  $\text{TMSCl}$  on the 0-4 position, resulting in the production of the 6-0-benzyl ether. This difference is caused by the higher sensitivity of the p-methoxybenzylidene group, resulting from the electron-donating effect of its methoxyl group.

## 2-1-2 Synthesis of 3-0-benzyl derivatives

The synthesis of aglycons 40 and 41 was carried out by



methods similar to those used for the synthesis of aglycons 17 and 39, respectively, as shown in Fig.8. The starting material 44 was prepared by the conventional method.<sup>38),105)</sup>

Interestingly, the acetyl group at the O-2 position of compound 25 was found to be stable toward methanolysis in the presence of a catalytic amount of sodium methoxide (0.1 eq) in methanol at 0°C for one hour; 2,2,2-trichloroethyl 2-O-acetyl-3-O-benzyl-β-D-glucoside was obtained in 95% yield. A similar result was reported by Takeo and Tei<sup>98)</sup> in the methanolysis of benzyl 2,4,6-tri-O-acetyl-3-O-benzyl-α-D-glucopyranoside. Under the above conditions, both acetyl groups of 2,2,2-trichloroethyl 2,4-di-O-acetyl-3,6-di-O-benzyl-β-D-glucopyranoside (which was obtained from compound 41) were also stable; the extent of the deacetylation was only about 5% based on TLC. On the other hand, compound 12 and 2,2,2-trichloroethyl 2,3,4-tri-O-acetyl-6-O-benzyl-β-D-glucopyranoside (which was obtained from compound 39) were readily deacetylated to give compound 13 and 2,2,2-trichloroethyl 6-O-benzyl-β-D-glucopyranoside, respectively, in a quantitative yield.

These experimental results indicate that the benzyl group introduced into the O-3 position makes the acetyl groups at the O-2 and O-4 positions (i.e. adjacent to the benzyl group) stable toward such a mild methanolysis, but the benzyl group introduced into the O-6 position does not exert such a stabilization effect for the acetyl group. However, with compound 25 it seems that the acetyl group at the O-6 position is first subjected to methanolysis, thus




Figure 8. Synthetic route for compounds **40** and **41** from compound **44**.



compound 49 with  $\text{NaCNBH}_3$ - $\text{TMSCl}$  in acetonitrile gave the desired 6-O-benzyl ether 41 in 60% yield.

### 2-1-3 Glycosylation of 3-O-acetyl and 3-O-benzyl derivatives

The glycosylation reactions between  $\alpha$ -imidate 34 and aglycons 17,39-41 were conducted under the reaction conditions reported by Kawada *et al.*<sup>44)</sup>

A solution of  $\alpha$ -imidate 34 (2.0 eq) and the respective aglycon (1.0 eq) dissolved in anhydrous methylene chloride was added dropwise to a suspension of molecular sieves 4A in anhydrous methylene chloride below  $-70^\circ\text{C}$ . Boron trifluoride etherate (0.2 eq) in anhydrous methylene chloride was added dropwise to the above suspension below  $-70^\circ\text{C}$ . After the prescribed time (Table 1), the reaction mixture was worked up and the product was purified by preparative TLC. The configuration of the glucosidic center produced was determined by  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR spectroscopies. The results are summarized in Table 1.

At low temperature in the presence of boron trifluoride catalysis, highly stereoselective conversion of the  $\alpha$ -imidate to the  $\beta$ -glycoside generally proceeds smoothly with Walden inversion at the glycosidic center.<sup>83),84)</sup> Different from this general result, the glycosylation of aglycons 17 and 39 (with an acetyl group at the O-3 position) with  $\alpha$ -imidate 34 gave preferentially the  $\alpha$ -glycoside ( $\alpha/\beta$  ratio = about 1.8 and 2.5, respectively). Furthermore, the total yield was low (about 50% yield) and about 50% of the

Table 1. Results of glycosylation reactions between  $\alpha$ -imidate **34** and aglycons **17**, **39-41**

Aglycon	R <sub>1</sub> <sup>*</sup>	R <sub>2</sub> <sup>**</sup>	Reaction time (h)	Products	Yield (%)	
					$\beta$	$\alpha$
<b>17</b>	-Ac	-Ac	2	<b>37</b> ( $\beta$ ) <b>38</b> ( $\alpha$ )	16.4	29.0
<b>39</b>	-Ac	-Bzl	2	<b>51</b> ( $\beta$ ) <b>52</b> ( $\alpha$ )	13.0	33.0
<b>40</b>	-Bzl	-Ac	1	<b>53</b> ( $\beta$ )	94.2	--
<b>41</b>	-Bzl	-Bzl	1	<b>54</b> ( $\beta$ )	96.0	--

\* R<sub>1</sub> = 3-O substituent group


\*\* R<sub>2</sub> = 6-O substituent group

starting aglycon was recovered. Contrary to these results, the glycosylation of aglycons **40** and **41** (with a benzyl group at the 0-3 position) with  $\alpha$ -imidate **34** gave only the  $\beta$ -glycoside in extremely high yield. The effect of the benzyl group introduced into the 0-6 position on the  $\alpha/\beta$  ratio seems to be small. These results coincide well with those reported by Sinay<sup>87)</sup>. Thus, it is clear in the present experiments that the benzyl substituent at the 0-3 position of the aglycon is indispensable to obtain a  $\beta$ -D-(1  $\rightarrow$  4) linked glycoside with excellent yield.

The differences in the reactions that occur with the different aglycons may be explained as follows. The electron-donating benzyl group introduced into the 0-3 position (aglycons **40** and **41**) seems to increase the electron

density of the hydroxyl group at the C-4 position, i.e., the hydroxyl group becomes more nucleophilic. As the result, substitution by the aglycons (40 and 41) via a complete  $S_N2$  mechanism at the glucosidic center becomes favorable to afford only the  $\beta$ -glycoside. In contrast, the electron-withdrawing acetyl group at the O-3 position reduces the nucleophilicity of the hydroxyl group attached to the C-4 position of aglycon. Therefore, substitution by the aglycons (17 and 39) is thought to proceed with an  $S_N1$ -like mechanism, i.e., elimination of the trichloroacetimidoyl group in  $\alpha$ -imidate 34 is followed by a reaction between the aglycons (17 and 39) and the resulting carbonium ion (or oxonium ion) intermediate to afford preferentially the  $\alpha$ -glycoside, which is thermodynamically more stable than the  $\beta$ -glycoside.

The stabilization of the acetyl group in compound 25 and the results obtained from the above glycosylation experiments may be explained by the same electronic factor.



The benzyl group introduced into the O-3 position acts to increase the electron densities of the adjacent oxygen atoms (O-2 and O-4) to contribute to the stabilization of the acetyl groups at these positions with respect to mild methanolysis. At the same time, an increase in the nucleophilicity of the adjacent hydroxyl groups favors a complete  $S_N2$  reaction in the glycosylation with  $\alpha$ -imidate **34**. If such reasoning is correct, the presence of a benzyl (or other alkyl) group at the O-3 position of the aglycon will be important for the synthesis of a  $\beta$ -(1 $\rightarrow$ 2) glycoside and the presence of a benzyl (or other alkyl) group at the O-2 and/or the O-4 position of the aglycon will be important for the synthesis of a  $\beta$ -(1 $\rightarrow$ 3) glycoside.

It is clear that in the present work the  $\alpha/\beta$  ratio produced by the glycosylation largely depends on the protective groups of the aglycons, even if the imidate method is used.

## 2-2 Substituent-effect of other alkyl and acyl groups


The substituent-effect of 3-O-benzyl group may be explained by the inductive effect of the electron-donating benzyl group, but the possible participation of  $\pi$ -electrons on the benzene ring such as  $\pi$ -complex formation,<sup>14)</sup> a hydrogen-bond formation<sup>66)</sup> and so on cannot be ruled out. The former part of this section describes the effects of the  $\pi$ -electron of the benzyl group on the glycosylation reaction.

On the other hand, the removal of acyl group is easier

than that of benzyl group, particularly in the case of a high molecular weight compound. Therefore, the acyl group is more favorable as a persistent protective R-group. Then, the effect of the acyl groups other than acetyl group on the glycosylation reaction are also described in the latter part of this section.

### 2-2-1 Substituent-effect of 3-O-alkyl group

Three aglycons 55-57 were selected, for the glycosylation reaction with  $\alpha$ -imidate 34, in order to investigate the influence of the  $\pi$ -electron of the 3-O-benzyl substituent group on glycosylation reaction. Propyl, allyl and propargyl groups were chosen as typical models of single, double and triple bonds, respectively. The propargyl group can be converted into propyl and allyl groups by catalytic hydrogenation.



### Synthesis of aglycons

The synthetic route for compounds 55-57 from 1,2,4,6-tetra-O-acetyl-3-O-propargyl- $\beta$ -D-glucopyranose (58) is shown in Fig. 9. The starting material 58 was prepared via 1,2;5,6-di-O-isopropylidene- $\alpha$ -D-glucofuranose.<sup>105)</sup> The






Figure 9. Synthetic route for compounds 55-57 from compound 58.

synthesis of compound 57 was carried out by the methods similar to those used for the synthesis of compound 40.

Compounds 55 and 56 were obtained from compound 57 by catalytic hydrogenation in methanol using palladium carbon (Pd-C) in about 60% yield and using Lindlar catalyst (Pd-BaSO<sub>4</sub>)<sup>11)</sup> in about 30% yield, respectively. The reaction time should be about 30 min, because 2,2,2-trichloroethyl group was found to be unstable under the reaction conditions.

### Glycosylation

The glycosylation reaction between  $\alpha$ -imidate 34 and aglycons 55-57 was conducted under the same reaction conditions described in the previous section (Reaction time was one hour.). The product was purified by preparative TLC. The configuration of the anomeric center was determined by <sup>1</sup>H-NMR and <sup>13</sup>C-NMR spectroscopies.

The results are summarized in Table 2, which also contains the substituent constants ( $\sigma_1$ ) of R-groups<sup>7)</sup> and pKa-values of R-OH predicted by the method of Perrin et al.<sup>75)</sup>

As expected, three aglycons 55-57 gave only the  $\beta$ -glycoside in good yields, and 3-O-benzyl derivative 40 gave  $\beta$ -glycoside in an excellent yield as described in the previous section. Generally, the electron-donating ability of the C-3 functional group (R-) is evaluated by the substituent constant ( $\sigma_1$ ) and the pKa-value of the corresponding alcohol (R-OH).<sup>7),75)</sup> The yields of  $\beta$ -glycosides obtained by the reaction between glycon 34 and

Table 2. Results of glycosylation reactions between  $\alpha$ -imidate **34** and aglycons **40**, **55-57**

Aglycon	-R*	Product	Yield (%)	$\sigma_I^{**}$	pKa***
<b>55</b>	$-\text{CH}_2\text{CH}_2\text{CH}_3$	<b>63</b> ( $\beta$ )	79.3	-0.01	16.1
<b>56</b>	$-\text{CH}_2\text{CH}=\text{CH}_2$	<b>64</b> ( $\beta$ )	72.5	0.02	15.9
<b>57</b>	$-\text{CH}_2\text{C}\equiv\text{CH}$	<b>65</b> ( $\beta$ )	71.7	0.14	14.7
<b>40</b>	$-\text{CH}_2\text{C}_6\text{H}_5$	<b>53</b> ( $\beta$ )	94.2	0.03	15.5

\* -R = 3-O substituent group

\*\*  $\sigma_I$  = substituent constant for -R

\*\*\* pKa = value of R-OH

aglycons **55-57** seem to increase with decreasing of the  $\sigma_I$ -constant and with increasing of pKa-values. The yield of  $\beta$ -glycoside apparently is affected only by the inductive effect of the C-3 functional groups, not by the effect of the  $\pi$ -electrons of the C-3 functional groups.

Although  $\sigma_I$ -constant of benzyl group (0.03) is the similar to that of allyl group (0.02), the yield of  $\beta$ -glycoside from aglycon **40** is different from that from aglycon **56**. If the yield of the  $\beta$ -glycoside is affected only by the inductive effect of the C-3 functional group, both yields from aglycon **40** and **56** should be almost the same. Therefore, the excellent yield in the case of aglycon **40** should be affected not only by the inductive effect, but also by the other effect, i.e. by some through-space


interaction<sup>7)</sup> such as  $\pi$ -complex or a hydrogen bond formation between 3-O-benzyl and C-4 hydroxyl groups.

It is concluded that the benzyl group has a special function different from the other alkyl groups on  $\beta$ -glycosylation reaction. Benzyl group is especially useful for the protection of C-3 hydroxyl group of aglycon.

### 2-2-2 Substituent-effect of the 3-O-acyl group

Carbamoyl and pivaloyl groups which are more electron-donating than acetyl group may be more useful for protection of the starting glucose derivative 1. In fact, carbamoyl group has been used for the protection of the starting glucose derivative in the synthesis of cellulose-like polymer by Husemann and Müller<sup>39)</sup> and by Hirano.<sup>37)</sup> And pivaloyl group has been conveniently used for the  $\beta$ -glycosylation instead of acetyl group recently.<sup>50)</sup> On the other hand, the use of electron-withdrawing protective group of aglycon at O-3 position may be favorable for the formation of  $\alpha$ -glycoside; namely the more electron-withdrawing acyl group such as monochloroacetyl group may be useful for the synthesis of malto-oligosaccharides.

Then, four aglycons 66-69 were selected for the present investigation. Carbamoyl derivative 65 has poor solubility for dichloromethane used as a solvent in the glycosylation reaction. Since the aglycon 66 requires tedious handling, the aglycon 67 was also prepared as a carbamoyl derivative.



### Synthesis of aglycons

The synthetic route for compounds **66-69** from 2,2,2-trichloroethyl  $\beta$ -D-glucopyranoside (**13**) is shown in Fig. 10. Since carbamoyl and pivaloyl groups seem to migrate less readily than acetyl group,<sup>99)</sup> the C-6 hydroxyl group was acylated selectively to lead to compounds **66-68**. The selective carbamoylation<sup>36), 68)</sup> of compound **72** was carried out by the treatment with phenyl isocyanate in anhydrous pyridine at 50°C to give compound **66** in 58.9% yield. In this reaction, anhydrous pyridine was used to prevent the formation of by-products. The selective acetylation<sup>86), 100)</sup> of compound **72** was performed by the treatment with acetyl chloride / 2,6-lutidine in ethyl acetate at 0°C to afford compound **67** in 64.5% yield. The selective pivaloylation<sup>22), 58)</sup> of compound **73** was conducted by the treatment with pivaloyl chloride / pyridine in chloroform at 50°C to form compound **68** in 64.3% yield. The synthesis of compound **69** was conducted by the methods similar to those used for the synthesis of compound **17**.

### Glycosylation

The glycosylation reactions between  $\alpha$ -imide **34** and




Figure 10. Synthetic route for compounds **66-69** from compound **13**.

aglycons 66-69 were performed under the same reaction conditions as those described in the previous section (Section 2-1-3). The results are summarized in Table 3.

Contrary to the expectation from its reaction mechanism for imidate method, four aglycons 66-69 gave both  $\beta$ -glycoside and  $\alpha$ -glycoside. The  $S_N1$  reaction mechanism seems to participate to some extent in all these reactions and the extent of the participation depends on the protective groups used. The total yields of glycoside increased in the order of pivaloyl, carbamoyl, acetyl and monochloroacetyl groups. This result shows that the aglycon with more electron-donating acyl groups is more effective than that with more electron-withdrawing acyl group for the glycosylation reaction. Only carbamoyl derivatives 66 and 67 gave

Table 3. Results of glycosylation reactions between  $\alpha$ -imidate 34 and aglycons 17, 66-69

Aglycon	$-R_1^*$	$-R_2^{**}$	Products	Yield (%)		
				$\beta$	$\alpha$	total
66	-Car	-Car	75( $\beta$ ) 76( $\alpha$ )	35.3	20.3	55.6
67	-Car	-Ac	77( $\beta$ ) 78( $\alpha$ )	57.0	7.7	64.7
68	-Piv	-Piv	79( $\beta$ ) 80( $\alpha$ )	23.8	55.6	79.4
69	-MCA	-MCA	81( $\beta$ ) 82( $\alpha$ )	3.2	16.2	19.4
17	-Ac	-Ac	37( $\beta$ ) 38( $\alpha$ )	16.4	29.0	45.4

\*  $-R_1$  = 2-O and 3-O substituent groups

\*\*  $-R_2$  = 6-O substituent group

preferentially the  $\beta$ -glycosides, though the other acyl derivatives afforded preferentially the  $\alpha$ -glycosides.

Consequently, the effect of the C-3 substituent group of the aglycon on glycosylation by imidate method can be explained by the inductive effect. The C-3 electron-donating ether group seems to increase the electron density and nucleophilicity of the hydroxyl group at the C-4 position, and a complete  $S_N2$  reaction proceeds to afford only the  $\beta$ -glycoside. Especially, benzyl group with aromatic  $\pi$ -electron seems to have a special function such as through-space interaction. In contrast, on the glycosylation using aglycon with C-3 electron-withdrawing ester group,  $S_N1$  type reaction proceeds to give a mixture of  $\beta$ -glycoside and  $\alpha$ -glycoside. Thus, mainly, the inductive effect of the C-3 substituent of the aglycon is important factor to decide the configuration of the product in the glycosylation by imidate method. Further investigation of the interesting effect of aromatic  $\pi$ -electron of benzyl group may be needed.

In the cello-oligosaccharides synthesis, the starting material 1 should be protected with benzyl group at O-3 position.

## 2-3 Summary

In this chapter, substituent-effect of aglycon on glycosylation reaction was investigated.

A series of aglycons with different combinations of benzyl and acetyl groups at the O-3 and O-6 positions were



synthesized with good yield and glycosylation reactions with imidate 34 were carried out to study the influence of substituent groups on the reactivity of C-4 hydroxyl group of aglycon (Section 2-1). The aglycon with benzyl group at O-3 position afforded only the  $\beta$ -glycoside in extremely high yield, but the aglycon with acetyl group at O-3 position gave preferentially the  $\alpha$ -glycoside in about 30% yield (Table 1). The result revealed that the benzyl substituent at O-3 position of aglycon was indispensable to obtain only the  $\beta$ -D-(1 $\rightarrow$ 4) linked glycoside with excellent yield. The effect of 6-O-benzyl group on glycosylation reaction seemed to be small.

Although substituent-effect of 3-O-benzyl group can be explained by the inductive effect of the electron-donating benzyl group, the possible participation of  $\pi$ -electrons on benzene ring may also be important. Then, the influence of the  $\pi$ -electrons of the benzyl group on glycosylation reaction was studied (Section 2-2-1). As expected, all aglycons with 3-O-alkyl groups other than 3-O-benzyl group gave only the  $\beta$ -glycoside in good yields, and aglycon with 3-O-benzyl group afforded  $\beta$ -glycoside in the highest yield (Table 2). the results suggested that the aromatic  $\pi$ -electrons of 3-O-benzyl group, different from olefinic or acetylenic  $\pi$ -electrons, have a special function for increasing the yield of  $\beta$ -glycoside.

In addition, the effect of acyl group other than acetyl group was also investigated (Section 2-2-2). However, all aglycons with 3-O-acyl groups gave a mixture of  $\beta$ - and  $\alpha$ -

glycosides (Table 3). Among the acyl groups investigated, carbamoyl group was effective for  $\beta$ -glycoside formation. Thus, it was concluded that benzyl group was especially useful for the protection of C-3 hydroxyl group of the aglycon.

## Chapter 3 Substituent-effect of glycon on glycosylation reaction


### 3-1 Substituent-effect of the $\alpha$ -imide on glycosylation reaction

The benzyl group is necessary as the persistent group at the O-3 position of starting material 1 for the glycosylation reaction. However, the remaining protective groups at the O-2, O-4 and O-6 positions in the glycon (i.e., 3-O-benzyl- $\alpha$ -imide) will affect on glycosylation reaction. In this section, therefore, the stability of 3-O-benzyl- $\alpha$ -imides and the influence of the protective groups in the imides on the reactivity in the glycosylation reaction were investigated.

In general an unstable material has a high reactivity. For example,  $\alpha$ -imide 34 was too unstable to decompose on silica gel during the purification<sup>44)</sup>, but gave  $\beta$ -glycoside in excellent yield. However, the unstability of compound 34 is inconvenient for handling, especially in the case of large scale preparation. Then, the favorable  $\alpha$ -imide used for the cello-oligosaccharides synthesis should be highly reactive, but should be nevertheless stable on silica gel. In the present study, typical electron-donating p-methoxybenzyl and electron-withdrawing monochloroacetyl groups as a temporary protective group at O-4 position, and benzyl, acetyl and pivaloyl groups as persistent protective groups at O-2 and O-6 positions were initially selected,

respectively. Thus, six imidates (glycons) 34,83-87 were prepared to investigate their stability and reactivity on the glycosylation reaction.

On the other hand, Boeckel *et al.*<sup>3),4),5)</sup> reported that the effects exerted by O-4 and O-6 substituent groups are reversed, though these substituent groups are separated by the same number of single bonds from the anomeric center. They demonstrated that a combination of the electron-withdrawing acyl group at the O-4 position and the electron-donating alkyl group at the O-6 position facilitate the formation of the  $\beta$ -glycoside. Paulsen<sup>73),74)</sup> reported that benzyl compounds are always more reactive than the acetylated or benzoylated derivatives, the position of benzyl groups being less important than their number. For the clarification of these effects, three imidates (glycons) 88-90 were also prepared.



Compound 40 was selected as an aglycon, since this compound gave  $\beta$ -glycoside in a high yield and was relatively easily prepared from compound 44 as described in the previous chapter (Chapter 2, Section 2-1).

### 3-1-1 Synthesis of the $\alpha$ -imidates

#### Synthesis of $\alpha$ -imidates [I]

Imidate 34 was prepared by the method reported by Kawada et al.<sup>44)</sup> The synthetic route for imidates 83, 84, 88 and 90 from trichloroethyl 2,4,6-tri-O-acetyl-3-O-benzyl- $\beta$ -D-glucopyranoside (25) is shown in Fig. 11. The key reaction in the synthesis of compound 83 and 84 is the reductive cleavage of 4,6-O-p-methoxybenzylidene glucopyranoside 93 and 94 (Fig. 11, step f).

Johansson and Samuelsson<sup>42)</sup> reported that the reductive ring-opening of methyl 2,3-di-O-benzyl-4,6-O-p-methoxybenzylidene- $\alpha$ -D-glucopyranoside with  $\text{NaCNBH}_4$ -TMSCl in acetonitrile gave the 4-O-p-methoxybenzyl and 6-O-p-methoxybenzyl derivatives in 76% and 12% yields, respectively. However, different from the results of Johansson and Samuelsson<sup>42)</sup>, the acetyl derivative 93 afforded 4-O-p-methoxybenzyl ether 95 and 6-O-p-methoxybenzyl ether 96 in 38.5% and 30.0% yields by the treatment of  $\text{NaCNBH}_4$ -TMSCl in  $\text{CH}_3\text{CN}$ . On the other hand, the pivaloyl derivative 94 gave only compound 98 in 60.3% yield. The regioselectivity of the reductive cleavage of 4,6-O-p-methoxybenzylidene derivative seems to depend upon the electronic factor resulting from the substituent groups




Figure 11. Synthetic route for **83**, **84**, **88** and **90** from compound **25**

at the O-1 or O-2 position. Thus, the substituent-effect on the reductive ring-opening reaction should be more studied.

Imidate 83 was prepared from compound 95 by acetylation, detrichloroethylation and imidoylation. Since compound 96 was not obtained by the selective cleavage of compound 94, imidate 84 was prepared from compound 99 via compounds 100 and 101. Imidates 88 and 90 were synthesized from compounds 97 and 25, respectively.

#### Synthesis of $\alpha$ -imidates [2]

The synthetic route for imidates 85,86,87 and 89 is shown in Fig. 12. These imidates 85,86 and 87 were synthesized from compounds 35, 40 and 113 by monochloroacetylation, deprotection at the C-1 position (deallylation or detrichloroethylation), and imidoylation, respectively.

Imidate 89 was prepared from allyl 2,3-di-O-benzyl-4-O-p-methoxybenzyl- $\beta$ -D-glucopyranoside (106).<sup>43)</sup> The selective deallylation of compounds 108 and 111 with acyl group was done by the oxidative cleavage with selenium (IV) dioxide in acetic acid / dioxane<sup>45)</sup> to give compounds 109 and 112 in about 60 % yields, respectively.

#### **3-1-2 Stability of the $\alpha$ -imidates and their glycosylation**

The stability of the respective imidate was evaluated by the stability on the purification by preparative TLC (layer thickness 2mm, Kiesel gel 60 F<sub>254</sub> Merk) after imidoylation. The reactivities of imidates 34,83-90 were evaluated by the glycosylation reactions with aglycon 40




Figure 12. Synthetic route for compounds **85-87** and **89** from compounds **40**, **106** and **113**.



conducted under the similar reaction conditions as described previously; each imide (1.0 eq) was reacted with aglycon 40 (2.0 eq) in the presence of a catalytic amount of  $\text{BF}_3\text{-Et}_2\text{O}$  (0.2 eq) below  $-70^\circ\text{C}$  for one hour. The product was purified by a preparative TLC and identified by  $^1\text{H-NMR}$  analysis. The results are summarized in Table 4. In the case of imides 86-90, the corresponding  $\beta$ -glycoside was not isolated by TLC; the yield was less than 1 % and only hydrolysis of the imide was observed.

The introduction of the electron-withdrawing acyl group into  $\alpha$ -imide was found to contribute to the stabilization

Table 4. Results of the stability and reactivity of the  $\alpha$ -imides 34, 83-90

$\alpha$ -Imide	$-\text{R}_1^*$	$-\text{R}_2^{**}$	$-\text{R}_3^{***}$	Stability	Products	Yield (%)
34	-Bzl	-pMBzl	-Bzl	x	53 ( $\beta$ )	82.0
83	-Ac	-pMBzl	-Ac	o	118 ( $\beta$ )	60.0
84	-Piv	-pMBzl	-Piv	o	119 ( $\beta$ )	57.8
85	-Bzl	-MCA	-Bzl	o	120( $\beta$ )121( $\alpha$ )	13.5( $\beta$ )3.0( $\alpha$ )
86	-Ac	-MCA	-Ac	o	--	--
87	-Piv	-MCA	-Piv	o	--	--
88	-Ac	-Ac	-pMBzl	o	--	--
89	-Bzl	-Ac	-Ac	o	--	--
90	-Ac	-Ac	-Ac	o	--	--

\*  $-\text{R}_1$  = 2-O substituent group

\*\*  $-\text{R}_2$  = 4-O substituent group

\*\*\*  $-\text{R}_3$  = 6-O substituent group

of imidate, and to decrease the yield of  $\beta$ -glycoside. The replacement of the two benzyl groups at O-2 and O-6 positions of imidate 34 by acyl groups results in the stabilization of the imidate, and the  $\beta$ -glycosides were obtained in about 60% yields (imidates 83 and 84). Generally, pivaloyl derivative is more effective for the  $\beta$ -glycosylation than the corresponding acetyl derivative,<sup>50)</sup> but these two acyl derivatives, imidates 83 and 84, have almost the same reactivity. Then, the replacement of 4-O-p-methoxybenzyl group of the imidate 34 by the electron-withdrawing monochloroacetyl group extremely decreases the yield of  $\beta$ -glycoside (imidate 85). Thus, the electron-donating (p-methoxybenzyl) group at O-4 position of  $\alpha$ -imidate is very important for the formation of  $\beta$ -glycoside. As expected from these results, the stable imidates 86 and 87 did not afford any  $\beta$ -glycoside.

The replacement with each other between p-methoxybenzyl group at O-4 position and acetyl group at O-6 position of imidate 83 results in the formation of imidate 88. Contrary to the demonstration by Boeckel et al.,<sup>3),4),5)</sup> imidate 88 with the electron-withdrawing (acetyl) group at O-4 position and electron-donating (p-methoxybenzyl) group at O-6 position also did not afford  $\beta$ -glycoside. The result indicates that their demonstration is inapplicable to the present imidate method. Each of imidates 83, 88 and 89 has two electron-donating (benzyl and/or p-methoxybenzyl) groups, but only imidate 83 gives the  $\beta$ -glycoside in a moderate yield. The comparison of these results indicate

that the most effective group for the synthesis of  $\beta$ -glycoside is the ether group at C-4 position of the  $\alpha$ -imide, and that C-2 and C-6 ether groups are not so effective.

Imide 90 did not give  $\beta$ -glycoside. The additional introduction of benzyl group into O-2 position (imide 89) or O-6 position (imide 88) is not also effective, but the introduction of benzyl groups both into O-2 and O-6 positions results in the production of  $\beta$ -glycoside even if the strong electron-withdrawing monochloroacetyl group is introduced into O-4 position of the imide (imide 85). Thus, both benzyl groups at O-2 and O-6 positions may synergistically effect for the glycoside formation.

Thus, it is concluded that the most effective group for the  $\beta$ -glycoside synthesis is an electron-donating ether group such as p-methoxybenzyl groups at O-4 position of  $\alpha$ -imide and that the other benzyl groups at O-2 and O-6 positions have synergistical effects.

It is interesting that both electron-donating groups at O-3 position of aglycon and at O-4 position of glycon are very effective for the  $\beta$ -glycosidic bond formation. Therefore, protective groups should be considered as "reaction directing groups", not simply as "blocking groups".


### 3-2 Conclusive selection of starting material for cello-oligosaccharides synthesis

In the synthetic design proposed (Chapter 1, Section 1-1), the starting material 1, which can be converted both into glycon and aglycon by simple operations, is required. This section describes the synthesis of actual starting material discussed in Chapter 2 and in the previous section of this chapter.

Among the  $\alpha$ -imidates 34, 83-90 investigated,  $\alpha$ -imidates 83 and 84 are the most suitable in the present cellobiosaccharides synthesis because of its stability and reactivity. In addition, the product synthesized from the combination of  $\alpha$ -imide 83 and aglycon 40 may be used for the preparation of regiospecific substituted cellulose derivatives.

However, there is a problem in the synthesis of compounds 83 or 84, i.e.; the regioselectivity in the reductive cleavage of 4,6-O-p-methoxybenzylidene acetal. Kawada *et al.*<sup>43)</sup> reported that the reductive ring-opening of allyl-2,3-di-O-benzyl-4,6-O-p-methoxybenzylidene- $\beta$ -D-glucopyranoside with lithium aluminium hydride ( $\text{LiAlH}_4$ ) - aluminium chloride ( $\text{AlCl}_3$ ) gave the 4-O-p-methoxybenzyl ether derivative in 96.1% yield. On the other hand, Liptak<sup>57)</sup> reported that regioselectivity in the reductive cleavage of benzylidene acetal with  $\text{LiAlH}_4$ - $\text{AlCl}_3$  is associated with the steric hindrance exerted by the C-3 substituent. However, 2,2,2-trichloroethyl group is unstable under the reaction conditions of the reductive cleavage with  $\text{LiAlH}_4$ - $\text{AlCl}_3$ .<sup>28), 69)</sup> Then, allyl group was selected as a temporary X-group instead of 2,2,2-trichloroethyl group.

Allyl group is stable under a variety of conditions and can be selectively removed easily by the methods such as mild two-step reactions (isomerization of the double bond and subsequent acid hydrolysis of the resultant enol ether), oxidative cleavage with selenium (IV) dioxide.<sup>26),30),45)</sup> The latter reaction conditions are applicable to the acetyl derivatives. For the protective group pattern except C-1 position, the protective group patterns of compounds 83 and 84 were chosen. The p-methoxybenzyl group selected as a temporary Y-group can be removed oxidatively by the use of ammonium cerium (IV) nitrate (CAN)<sup>42)</sup> or 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ)<sup>72)</sup> without any influence of persistent acetyl, pivaloyl and benzyl groups. Therefore, allyl 2,6-di-O-acetyl-3-O-benzyl-4-O-p-methoxybenzyl-β-D-glucopyranoside (122) and allyl 2,6-di-O-pivaloyl-3-O-benzyl-4-O-p-methoxybenzyl-β-D-glucopyranoside (123) were selected as starting material 1 in the present cello-oligosaccharides synthesis.



The synthetic route for compounds 122 and 123 from compound 44 is shown in Fig. 13. The key reaction in this




Figure 13. Synthetic route for compounds 122 and 123 from compound 44

route is  $\beta$ -allylglycosylation (step a) and reductive ring-opening reaction (step d).

For the synthesis of  $\beta$ -allylglucopyranoside, a Koenigs-Knorr method in which acylglucosyl halide is reacted with allyl alcohol in the presence of a suitable catalyst such as silver carbonate<sup>21)</sup> or mercuric cyanide<sup>53)</sup> was tried. However, this method requires the additional halogenation step before glycosylation. Furthermore, 3-O-benzyl group of compound 44 was found to be partly removed under the halogenation reaction conditions. Then, the modified Magnusson's method<sup>13),59)</sup> was examined. Compound 44 was treated with allyl alcohol / boron trifluoride etherate to afford compound 124 in 77.4% yield. 3-O-Benzyl group was stable under this reaction conditions. In this reaction, it was found that allyl alcohol and boron trifluoride etherate should be used in excess (5.0 eq and 10.0 eq) for completion of the reaction. Allyl 2,3,4,6-tetra-O-acetyl- $\beta$ -D-glucopyranoside is also obtained from compound 11 in 70.5% yield. Thus, this procedure is a simple and highly stereoselective  $\beta$ -allylglycosylation method for the compound with benzyl ether.

The reductive ring-opening reaction of compound 126 using the  $\text{LiAlH}_4\text{-AlCl}_3$  gave 4-O-p-methoxybenzyl ether derivative 127 in 83.6 % yield. Compound 127 was converted to the target material 122 and 123 by acetylation and by pivaloylation, respectively, in a quantitative yield.

The starting material 122 was converted into glycon 102 and aglycon 128 by the removal of the ally and p-




Figure 14.

Synthetic route for compound **129** from compound **122**



methoxybenzyl groups, respectively. The allyl group was removed by the oxidation with selenium (IV) dioxide to produce glycon 102 in 61.0% yield. For the removal of p-methoxybenzyl group, compound 122 was treated with cerium (IV) ammonium nitrate to give aglycon 128 in 85.3% yield. Then,  $\alpha$ -imide 83 was obtained from compound 102. Glycosylation between  $\alpha$ -imide 83 and aglycon 128 gave the  $\beta$ -glycoside 129 in about 50% yield.

### 3-3 Summary

In Chapter 3, the influence of the substituent groups of the glycon ( $\alpha$ -imide) on their stability and reactivity was investigated (Section 3-1). As the results, it was found that the introduction of an electron-withdrawing acyl group into the  $\alpha$ -imide contributes to stabilization of the  $\alpha$ -imide, but decreases the yield of  $\beta$ -glycoside, and that the electron-donating ether group such as p-methoxybenzyl group introduced into O-4 position of  $\alpha$ -imide was very effective. The 2-O- and 6-O-alkyl groups are not so effective for the  $\beta$ -glycoside formation.

Finally, a favorable starting material for the synthesis of cello-oligosaccharides was designed on the basis of the results mentioned above (Section 3-2). Allyl 2,6-di-O-acetyl-3-O-benzyl-4-O-p-methoxybenzyl- $\beta$ -D-glucopyranoside (122) and allyl 2,6-di-O-pivaloyl-3-O-benzyl-4-O-p-methoxybenzyl- $\beta$ -D-glucopyranoside (123) were proposed as a conclusive starting material for the synthesis of cello-

oligosaccharides.

## CONCLUSIONS

There are two synthetic problems, regiospecific 1,4-bond formation and stereospecific  $\beta$ -linkage formation, have to be solved for the cello-oligosaccharides synthesis. The former problem will be solved by the use of a starting material which has special protective groups introduced by regiospecific control, the latter by the stereospecific  $\beta$ -glycosidic bond synthesis. The author investigated the synthetic methods for the starting materials and their glycosylation reaction in order to synthesize cello-oligosaccharides. The substituted glucose derivative 1 as a starting material should have three kinds of protective groups, X-, Y- and R-groups.

In Chapter 1, the synthesis of a starting material, in which acyl group is used as a R-group, and its glycosylation reaction for the regulation of regiospecificity were studied. The synthetic route for 2,3,6-tri-O-acetyl glucose derivatives from D-glucose (Fig. 3) was investigated at first. Here, two problems lay in tritylation at C-6 hydroxyl group (Fig. 3, step d) and acetyl group migration from O-4 to O-6 (Fig. 3, step g). The former problem was solved by the use of trityl chloride / triethylamine / dioxane system and the latter by the use of p-toluenesulfonic acid. 2,3,6-Tri-O-benzoyl glucose derivatives were also synthesized successfully. Then, the glycosylation between the acetyl derivative and  $\alpha$ -imidate 34 was examined. As a result, it was found that the yield and  $\alpha/\beta$  ratio of glycosides were

greatly affected by the protective group pattern.

In Chapter 2, a series of aglycons with different combinations of ether and ester groups at the C-3 and C-6 positions were successfully synthesized with good yields. Then, the glycosylation reactions with imidate 34 were conducted to investigate the influence of the substituent groups on the reactivity of the C-4 hydroxyl group of aglycons. The result indicated that the benzyl substituent at the O-3 position of the aglycon was indispensable to obtain only the  $\beta$ -D-(1 $\rightarrow$ 4) linked glycoside with excellent yield, by the effect of the  $\pi$ -electron of the benzene ring in addition to the inductive effect of the electron-donating benzyl group (Tables 1, 2 and 3).

In Chapter 3, the influence of the substituent groups of the glycon ( $\alpha$ -imidate) on their stability and reactivity was investigated. It was found that the introduction of an electron-withdrawing acyl group into the  $\alpha$ -imidate contributes to stabilization of the  $\alpha$ -imidate, but decreases the yield of  $\beta$ -glycoside, and that the electron-donating ether group such as p-methoxybenzyl group introduced into O-4 position of  $\alpha$ -imidate was very effective (Table 4). Thus, protective groups should be considered as "reaction directing groups" not simply as "blocking groups" from the present systematic investigation. The suitable selection and effective use of protective groups are very important for the synthesis of oligosaccharides and polysaccharides.

Finally, allyl 2,6-di-O-acetyl-3-O-benzyl-4-O-p-methoxybenzyl- $\beta$ -D-glucopyranoside (122) and allyl 2,6-di-O-pivaloyl-3-O-benzyl-4-O-p-methoxybenzyl- $\beta$ -D-glucopyranoside (123) were proposed as a conclusive starting material for the synthesis of cello-oligosaccharides.

## EXPERIMENTAL

### GENERAL

All melting points are uncorrected. Optical rotations were determined with a Jasco DIP-4 polarimeter for solutions in  $\text{CHCl}_3$  at  $28^\circ\text{C}$ , unless otherwise noted. UV spectra were recorded with a Shimadzu UV-365 spectrophotometer. IR spectra were recorded with a Shimadzu FT IR-4000 spectrophotometer.  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR spectra were taken with a Varian XL-200 FT-NMR (200 MHz) and a JEOL FX-NMR (90 MHz) spectrophotometers, respectively, with TMS internal standard in  $\text{CDCl}_3$ . Chemical shifts and coupling constants are given in  $\delta$ -value and Hz, respectively. Preparative TLC (hereafter "P-TLC") was done on silica gel plates (layer thickness 2 mm, Kieselgel 60 F  $_{254}$  Merck). Anhydrous  $\text{CH}_2\text{Cl}_2$ , anhydrous  $\text{CH}_3\text{CN}$ , anhydrous  $\text{CCl}_3\text{CH}_2\text{OH}$  and anhydrous pyridine were distilled from  $\text{P}_2\text{O}_5$ . Anhydrous dioxane was distilled from sodium metal. Anhydrous cyclohexane and anhydrous  $\text{Et}_2\text{O}$  were distilled from sodium hydride. The standard work-up included diluting with  $\text{EtOAc}$ , washing with a saturated  $\text{NaHCO}_3$  solution and with a saturated  $\text{NaCl}$  solution (hereafter "brine"), drying over  $\text{Na}_2\text{SO}_4$  and evaporating in vacuo.

## 1. EXPERIMENTAL (Chapter 1)

2,2,2-Trichloroethyl

2,3,4,6-tetra-O-acetyl- $\beta$ -D-glucopyranoside (12)

To a solution of glucosepentaacetate<sup>105)</sup> (11) (78 g, 0.2M) and 2,2,2-trichloroethanol (57.6ml, 0.6M) dissolved in  $\text{CH}_2\text{Cl}_2$  (100 ml), powdered molecular sieves 4A (20 g) was added. After stirring for 1 h at room temperature,  $\text{BF}_3\text{-Et}_2\text{O}$  (123ml, 1M) was added dropwise over a period of 1 h at  $0^\circ\text{C}$ . The reaction mixture was stirred for 15 h and then filtered. The residual molecular sieves was washed with EtOAc (200 ml). The combined filtrate and washings were worked-up by the standard method to give a colorless syrup which was crystallized from EtOH to afford colorless needles (60 g, 63.0% yield); m.p.  $144\text{--}145^\circ\text{C}$  (Lit.  $143\text{--}145^\circ\text{C}$ )<sup>59)</sup>,  $^1\text{H-NMR}$ : 2.02 (3H, s), 2.04 (3H, s), 2.07 (3H, s), 2.11 (3H, s), 3.72 (1H, ddd,  $J=10.0, 4.5, 2.4$ ,  $\text{C}_5\text{-H}$ ), 4.13 (1H, d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.14 (1H, dd,  $J=12.0, 2.5$ ,  $\text{C}_6\text{-H}_a$ ), 4.26 (1H, dd,  $J=12.0, 4.5$ ,  $\text{C}_6\text{-H}_b$ ), 4.80 (1H, d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.84 (1H, d,  $J=7.3$ ,  $\text{C}_1\text{-H}$ ), 5.10 (1H, broad t,  $J=10.0$ ,  $\text{C}_2\text{-H}$ ), 5.14 (1H, t,  $J=10.0$ ,  $\text{C}_4\text{-H}$ ), 5.24 (1H, broad t,  $J=10.0$ ,  $\text{C}_3\text{-H}$ ).

2,2,2-Trichloroethyl  $\beta$ -D-glucopyranoside (13)

To a suspension of 12 (4.79 g, 10 mM) in MeOH (70 ml), 28% NaOMe in MeOH (0.4 ml, 2 mM) was added at  $0^\circ\text{C}$ . The

reaction mixture was stirred for 1 h at room temperature. After treatment with Amberlite IR-120B (H<sup>+</sup>) ion-exchange resin, the resin was filtered off and washed with MeOH. The combined filtrate and washings were evaporated in vacuo to give a colorless syrup, which was triturated from EtOAc to afford colorless crystals (2.85 g, 91.4% yield); m.p. 148-149°C;  $[\alpha]_D -36.3^\circ$  (H<sub>2</sub>O). Anal. Calcd. for C<sub>8</sub>H<sub>13</sub>O<sub>6</sub>Cl<sub>3</sub>: C, 30.82; H, 4.17. Found: C, 30.91; H, 4.25.

2,2,2-Trichloroethyl 6-O-trityl- $\beta$ -D-glucopyranoside (14)

To a solution of 13 (2.49 g, 8 mM) and trityl chloride (2.68 g, 9.6 mM) in anhydrous dioxane, Et<sub>3</sub>N (5.6 ml) was added dropwise. The solution was kept at 100°C for 1 h, then cooled and diluted with EtOAc. The organic layer was washed with a saturated NH<sub>4</sub>Cl solution and with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated. The product was purified on a silica gel column (Wacogel C-100, 40 g, 12 cm x 3 cm) eluted with CH<sub>2</sub>Cl<sub>2</sub> to give 14 as a syrup (3.84 g, 86.9% yield), which was triturated from n-hexane to afford colorless crystals; m.p. 74-76°C;  $[\alpha]_D -17.3^\circ$  (dioxane), UV  $\lambda_{\text{max}}^{\text{EtOH}}$  nm( $\epsilon$ ): 260 (760), 254 (680). Anal. Calcd. for C<sub>27</sub>H<sub>27</sub>O<sub>6</sub>Cl<sub>3</sub>: C, 58.54; H, 4.88. Found: C, 58.82; H, 5.27.

2,2,2-Trichloroethyl 2,3,4-tri-O-acetyl-6-O-trityl- $\beta$ -D-glucopyranoside (15)

Compound 14 (1.66 g, 3 mM) and NaOAc (986 mg, 12 mM) were dissolved in Ac<sub>2</sub>O (40 ml). The reaction mixture was kept at 100°C for 1 h with vigorous stirring. The cooled



mixture was worked-up by the standard method. The residual  $\text{Ac}_2\text{O}$  was removed as the EtOH azeotrope. The viscous product was crystallized from EtOH / n-hexane to afford colorless crystals (1.84 g, 90.6% yield); m.p. 126-128°C;  $[\alpha]_D$  17.2°; UV  $\lambda_{\text{max}}^{\text{EtOH}}$  nm( $\epsilon$ ): 262 (790), 255 (740); IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 1760 (C=O), 1220 (C-O);  $^1\text{H-NMR}$ : 1.75 (3H, s), 2.01 (3H, s), 2.07 (3H, s), 3.11 (1H, dd,  $J=10.0$ , 4.5,  $\text{C}_6\text{-H}_a$ ), 3.31 (1H, dd,  $J=10.0, 2.5$ ,  $\text{C}_6\text{-H}_b$ ), 3.61 (1H, ddd,  $J=10.0, 4.5, 2.5$ ,  $\text{C}_5\text{-H}$ ), 4.23 (1H, d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.46 (1H, d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.84 (1H, d,  $J=7.5$ ,  $\text{C}_1\text{-H}$ ), 5.19 (1H, t,  $J=10.0$ ,  $\text{C}_2\text{-H}$ ), 5.24 (1H, t,  $J=10.0$ ,  $\text{C}_4\text{-H}$ ), 5.25 (1H, t,  $J=10.0$ ,  $\text{C}_3\text{-H}$ ), 7.34 (15H, broad, trityl). Anal. Calcd. for  $\text{C}_{33}\text{H}_{33}\text{O}_9\text{Cl}_3$ : C, 58.28; H, 4.86. Found: C, 58.27; H, 4.88.

**2,2,2-Trichloroethyl 2,3,6-tri-O-acetyl- $\beta$ -D-glucopyranoside (17)**

A solution of compound 15 (1.02 g, 1.5 mM) in  $\text{CH}_2\text{Cl}_2$  (20 ml) was stirred in the presence of p-toluenesulfonic acid (517 mg, 3 mM) at r.t. for 3 h. The solution was worked-up by the standard method. The product was purified on a silica gel column (Wacogel C-100, 20 g, 8 cm x 2.5 cm) eluted with  $\text{CH}_2\text{Cl}_2$  to give 17 as an oil, which was crystallized from EtOH / n-hexane to afford colorless crystals (569 mg, 86.8% yield); m.p. 136-138°C;  $[\alpha]_D$  -59.2°; IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 3550 (OH), 1740 (C=O), 1230 (C-O);  $^1\text{H-NMR}$ : 2.07 (3H, s), 2.11 (3H, s), 2.15 (3H, s), 3.54 (1H,

ddd,  $J=10.0$ , 4.5, 2.0,  $C_5-H$ ), 3.58 (1H, broad,  $C_4-H$ ), 4.15 (1H, d,  $J=12.0$ ,  $-OCH_2CCl_3$ ), 4.34 (1H, dd,  $J=12.0$ , 2.0,  $C_6-H_a$ ), 4.41 (1H, d,  $J=12.0$ ,  $-OCH_2CCl_3$ ), 4.51 (1H, dd,  $J=12.0$ , 4.5,  $C_6-H_b$ ), 4.83 (1H, d,  $J=7.5$ ,  $C_1-H$ ), 5.06 (1H, t,  $J=9.5$ ,  $C_2-H$ ), 5.10 (1H, t,  $J=10.0$ ,  $C_3-H$ ). Anal. Calcd. for  $C_{14}H_{19}O_9Cl_3$ : C, 38.40; H, 4.34. Found: C, 38.70; H, 4.29.

**2,2,2-Trichloroethyl 2,3,6-tri-O-acetyl-4-O-benzyl- $\beta$ -D-glucopyranoside (9)**

Compound 17 (44 mg, 0.1 mM) was dried over  $P_2O_5$  in a vacuum desiccator, and molecular sieves 4A powder (100 mg) was dried over night in a  $120^\circ C$  oven before use. A solution of 17 in anhydrous  $CH_2Cl_2$  (1 ml) was added to a stirred suspension of molecular sieves in anhydrous cyclohexane (1 ml). To this reaction mixture, benzyltrichloroacetimidate<sup>12)</sup> (56  $\mu$ l, 0.3 mM)<sup>41)</sup> was added. After stirring this mixture at room temperature for 30 min,  $CF_3SO_3H$  (8  $\mu$ l, 0.05 mM) was added. The reaction mixture was then kept at room temperature for an additional 5.5 h. The mixture was worked-up by the standard method to give a colorless oil. This oil was purified by P-TLC (EtOAc / *n*-hexane (1:4)) to give 9 (27.5 mg, 51.8% yield) which was crystallized from EtOH, m.p.  $156-158^\circ C$ ;  $[\alpha]_D -38.2^\circ$ ; UV  $\lambda_{max}^{dioxane}$  nm( $\epsilon$ ): 259(180), 253(140), 266(140); IR  $\nu_{max}^{KBr}$   $cm^{-1}$ : 1730 (C=O), 1230 (C-O);  $^1H$ -NMR: 2.00 (3H, s), 2.06 (3H, s), 2.08 (3H, s), 3.62 (1H, ddd,  $J=10.0$ , 4.5, 2.0,  $C_5-H$ ), 3.72 (1H, t,  $J=8.5$ ,  $C_4-H$ ), 4.13 (1H, d,  $J=12.0$ ,  $-OCH_2CCl_3$ ), 4.22 (1H, dd,  $J=12.0$ , 4.0,  $C_6-H_a$ ), 4.40 (1H, d,  $J=12.0$ ,  $-OCH_2CCl_3$ ), 4.44 (1H, dd,

$J=12.0, 2.0$ ,  $C_6-H_b$ ), 4.46 (1H, d,  $J=11.0$ , benzyl), 4.56 (1H, d,  $J=11.0$ , benzyl), 4.82 (1H, d,  $J=7.5$ ,  $C_1-H$ ), 5.03 (1H, t,  $J=8.5$ ,  $C_2-H$ ), 5.31 (1H, d,  $J=8.5$ ,  $C_3-H$ ), 7.30 (5H, phenyl).  
 Anal. Calcd. for  $C_{21}H_{25}O_9Cl_3$ : C, 47.78; H, 4.74. Found: C, 48.02; H, 4.73.

2,2,2-Trichloroethyl 3,4,6-tri-O-acetyl-2-O-benzyl- $\beta$ -D-glucopyranoside (24) and 2,2,2-trichloroethyl 2,4,6-tri-O-acetyl-3-O-benzyl- $\beta$ -D-glucopyranoside (25)

To a solution of the compound 17 (44 mg, 0.1 mM) and benzyl bromide (25.7 mg, 0.15 mM) dissolved in DMF (3 ml), NaH, 60% dispersion in mineral oil, (6 mg, 0.15 mM) was added at 0°C. After stirring for 30 min at 0°C, the reaction mixture was partitioned between  $Et_2O$  and brine. The  $Et_2O$  layer was dried over sodium sulfate and evaporated in vacuo to give a colorless oil which was purified by P-TLC (EtOAc / n-hexane (1:3)). The benzylated compound 9 (Rf-value = 0.5) was obtained in 17.0% yield (9 mg). Another separated fraction (Rf-value = 0.25, 8 mg) was further purified by P-TLC ( $CH_2Cl_2$  / n-hexane (2:1) three times) to afford compound 24 (Rf-value = 0.47, 4 mg, 7.6% yield) and compound 25 (Rf-value = 0.40, 2 mg, 3.8% yield).

Compound 24;  $^1H$ -NMR: 1.94 (3H, s), 2.02 (3H, s), 2.10 (3H, s), 3.53 (1H, dd,  $J=9.7, 7.7$ ,  $C_2-H$ ), 3.71 (1H, ddd,  $J=9.5, 4.2, 2.2$ ,  $C_5-H$ ), 4.14 (1H, dd,  $J=12.0, 2.2$ ,  $C_6-H_a$ ), 4.23 (1H, d,  $J=11.7$ ,  $-OCH_2CCl_3$ ), 4.30 (1H, dd,  $J=12.0, 4.2$ ,  $C_6-H_b$ ), 4.53 (1H, d,  $J=11.7$ ,  $-OCH_2CCl_3$ ), 4.68 (1H, d,

$J=11.8$ , benzyl), 4.80 (1H, d,  $J=7.7$ ,  $C_1$ -H), 4.95 (1H, d,  $J=11.8$ , benzyl), 4.99 (1H, t,  $J=9.5$ ,  $C_4$ -H), 5.17 (1H, t,  $J=9.5$ ,  $C_3$ -H), 7.26-7.44 (5H, m, phenyl).

Compound 25;  $^1\text{H-NMR}$ : 1.98 (3H, s), 2.04 (3H, s), 2.10 (3H, s), 3.63 (1H, ddd,  $J=9.5$ , 4.5, 2.0,  $C_5$ -H), 3.74 (1H, t,  $J=9.5$ ,  $C_3$ -H), 4.13 (1H, d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.14 (1H, dd,  $J=12.0$ , 2.0,  $C_6$ -H<sub>a</sub>), 4.22 (1H, dd,  $J=12.0$ , 4.5,  $C_6$ -H<sub>b</sub>), 4.40 (1H, d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.59 (1H, d,  $J=12.0$ , benzyl), 4.65 (1H, d,  $J=12.0$ , benzyl), 4.76 (1H, d,  $J=8.0$ ,  $C_1$ -H), 5.15 (1H, t,  $J=9.5$ ,  $C_4$ -H), 5.17 (1H, dd,  $J=9.5$ , 8.0,  $C_2$ -H), 7.20-7.42 (5H, m, phenyl).

2,2,2-Trichloroethyl

2,3,6-tri-O-acetyl-4-O-

methylthiomethyl- $\beta$ -D-glucopyranoside (26)

Compound 17 (44 mg, 0.1 mM) dissolved in dimethylsulfoxide (0.5 ml) was mixed with AcOH (0.5 ml) and  $\text{Ac}_2\text{O}$  (0.5 ml) and then heated at  $100^\circ\text{C}$  for 1 h. The mixture was diluted with EtOAc, washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated. The residual solvents were removed as the EtOH azeotrope. The product was crystallized from EtOH / n-hexane (31.7 mg, 63.4% yield), m.p.  $141-143^\circ\text{C}$ ;  $[\alpha]_D$   $4.0^\circ$ ; IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 1740 (C=O), 1230 (C-O);  $^1\text{H-NMR}$ : 2.05 (3H, s), 2.09 (3H, s), 2.13 (3H, s), 2.15 (3H, s), 3.60 (1H, ddd,  $J=10.0$ , 5.0, 2.5,  $C_5$ -H), 3.84 (1H, t,  $J=9.5$ ,  $C_4$ -H), 4.13 (1H, d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.35 (1H, dd,  $J=12.0$ , 4.0,  $C_6$ -H<sub>a</sub>), 4.38 (1H, d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.45 (1H, dd,  $J=12.0$ , 2.0,  $C_6$ -H<sub>b</sub>), 4.68 (2H, s,  $-\text{CH}_2\text{SCH}_3$ ), 4.82 (1H, d,  $J=9.0$ ,  $C_1$ -H), 5.02 (1H, dd,  $J=9.5$ , 7.5,  $C_2$ -H), 5.25

(1H, t, J=9.0, C<sub>3</sub>-H). Anal. Calcd. for C<sub>16</sub>H<sub>23</sub>O<sub>9</sub>Cl<sub>3</sub>S: C, 38.59; H, 4.62. Found: C, 38.73 ; H, 4.59.

2,2,2-Trichloroethyl 2,3,6-tri-O-acetyl-4-O-diphenylmethyl-β-D-glucopyranoside (27)

To a solution of 17 (65 mg, 0.15 mM) and benzhydrol (138 mg, 0.75 mM) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (1 ml), P<sub>2</sub>O<sub>5</sub> (about 300 mg) was added, and the mixture was stirred at room temperature for 15 min. The mixture was poured into a stirred suspension of EtOAc (5 ml) and NaHCO<sub>3</sub> (300 mg), and then diluted with H<sub>2</sub>O. The organic layer was washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated. The residue was purified by P-TLC (CH<sub>2</sub>Cl<sub>2</sub> / n-hexane (2:1)) to yield colorless crystals (60.4 mg, 66.7% yield); m.p. 154-155°C; [α]<sub>D</sub> 3.1°; UV λ<sub>max</sub><sup>dioxane</sup> nm(ε): 260(400), 254(350), 266(310); IR ν<sub>max</sub><sup>KBr</sup> cm<sup>-1</sup>: 1755 (C=O), 1235 (C-O); <sup>1</sup>H-NMR: 1.66 (3H, s), 1.94 (3H, s), 2.03 (3H, s), 3.66 (1H, ddd, J=10.0, 4.0, 2.0, C<sub>5</sub>-H), 3.84 (1H, dd, J=12.0, 4.0, C<sub>6</sub>-H<sub>a</sub>), 3.90 (1H, t, J=9.0, C<sub>4</sub>-H), 4.10 (1H, d, J=12.0, -OCH<sub>2</sub>CCl<sub>3</sub>), 4.23 (1H, dd, J=12.0, 2.0, C<sub>6</sub>-H<sub>b</sub>), 4.35 (1H, d, J=12.0, -OCH<sub>2</sub>CCl<sub>3</sub>), 4.84 (1H, d, J=8.0, C<sub>1</sub>-H), 4.96 (1H, dd, J=8.5, 7.5, C<sub>2</sub>-H), 5.35 (1H, t, J=9.0, C<sub>3</sub>-H), 5.50 (1H, s, -CH(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>), 7.24-7.29 (10H, m, phenyl). Anal. Calcd. for C<sub>27</sub>H<sub>29</sub>O<sub>9</sub>Cl<sub>3</sub>: C, 53.69 ; H, 4.80. Found: C, 53.69 ; H, 4.79.

2,2,2-Trichloroethyl 2,3,6-tri-O-acetyl-4-O-methoxymethyl-β-D-glucopyranoside (28)

To a solution of 17 (44 mg, 0.1 mM) and methyral (88  $\mu$ l, 1 mM) in anhydrous  $\text{CH}_2\text{Cl}_2$  (1 ml),  $\text{P}_2\text{O}_5$  (about 200 mg) was added. The reaction mixture was stirred at room temperature for 10 min. The mixture was worked-up in the same way as that of 27 to give a colorless oil. The product was crystallized from EtOH /n-hexane (37.2 mg, 76.6% yield); m.p. 89-90°C;  $[\alpha]_D$  -34.8°; IR  $\nu_{\text{max}}^{\text{kBr}}$   $\text{cm}^{-1}$  : 1745 (C=O) 1235 (C-O);  $^1\text{H-NMR}$  : 2.06 (3H, s), 2.08 (3H, s), 2.12 (3H, s), 3.32 (3H, s,  $-\text{OCH}_2\text{OCH}_3$ ), 3.61 (1H, ddd,  $J=9.5, 4.5, 2.0$ ,  $\text{C}_5\text{-H}$ ), 3.74 (1H, dd,  $J=9.5, 9.0$ ,  $\text{C}_4\text{-H}$ ), 4.13 (1H, d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.24 (1H, dd,  $J=12.0, 4.5$ ,  $\text{C}_6\text{-H}_a$ ), 4.39 (1H, d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.42 (1H, dd,  $J=12.0, 2.5$ ,  $\text{C}_6\text{-H}_b$ ), 4.58 (1H, d,  $J=6.5$ ,  $-\text{OCH}_2\text{OCH}_3$ ), 4.68 (1H, d,  $J=6.5$ ,  $-\text{OCH}_2\text{OCH}_3$ ), 4.83 (1H, d,  $J=7.5$ ,  $\text{C}_1\text{-H}$ ), 5.01 (1H, dd,  $J=9.5, 8.0$ ,  $\text{C}_2\text{-H}$ ), 5.22 (1H, dd,  $J=9.0, 8.5$ ,  $\text{C}_3\text{-H}$ ). Anal. Calcd. for  $\text{C}_{16}\text{H}_{23}\text{O}_{10}\text{Cl}_3$ : C, 39.88 ; H, 4.78. Found: C, 38.83 ; H, 4.63.

**2,2,2-Trichloroethyl 2,3,6-tri-O-benzoyl-4-O-methoxymethyl- $\beta$ -D-glucopyranoside (30)**

**Method A:** To a stirred solution of 28 (169 mg, 0.35 mM) in MeOH (4 ml), 28% NaOMe in MeOH (0.14 ml, 0.7 mM) was added at 0°C. The suspension was then stirred at room temperature for 1 h. The reaction mixture was worked-up in the same way as that of 13 to give a colorless oil. After the residue was dissolved in anhydrous pyridine (2 ml), benzoyl chloride (243  $\mu$ l, 2.1 mM) was added. The solution was heated at 50°C for 1.5 h, cooled, and then worked-up by the standard method. The residual pyridine was

removed as the ethanol azeotrope. The product was purified by P-TLC (EtOAc / *n*-hexane (1:4)) to afford **30** (149.8 mg, 63.5% overall yield).

Method B: A solution of **32** (50 mg, 0.08 mM) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (1 ml) was treated with methyral (88 μl, 1 mM) and P<sub>2</sub>O<sub>5</sub> (about 200mg), and stirred at room temperature for 10 min. The mixture was worked-up in the same way as that of **27** to give a colorless oil (42.2 mg, 77.5% yield). The product was crystallized from EtOH.

Compound **30**; m.p. 84-85°C; [ $\alpha$ ]<sub>D</sub> 51.5°; UV  $\lambda_{\text{max}}^{\text{dioxane}}$  nm  
( $\epsilon$ ): 275 (2680), 283 (2150); IR  $\nu_{\text{max}}^{\text{KBr}}$  cm<sup>-1</sup>: 1725 (C=O), 1275 (C-O); <sup>1</sup>H-NMR: 3.17 (3H, s, -OCH<sub>2</sub>OCH<sub>3</sub>), 3.93 (1H, ddd, J=9.5, 4.0, 2.0, C<sub>5</sub>-H), 4.11 (1H, t, J=9.5, C<sub>4</sub>-H), 4.16 (1H, d, J=12.0, -OCH<sub>2</sub>CCl<sub>3</sub>), 4.40 (1H, d, J=12.0, -OCH<sub>2</sub>CCl<sub>3</sub>), 4.56 (1H, dd, J=12.0, 4.0, C<sub>6</sub>-H<sub>a</sub>), 4.57 (1H, d, J=7.0, -OCH<sub>2</sub>OCH<sub>3</sub>), 4.69 (1H, d, J=7.0, -OCH<sub>2</sub>OCH<sub>3</sub>), 4.75 (1H, dd, J=12.0, 2.5, C<sub>6</sub>-H<sub>b</sub>), 5.09 (1H, d, J=8.0, C<sub>1</sub>-H), 5.48 (1H, dd, J=10.0, 7.5, C<sub>2</sub>-H), 5.74 (1H, dd, J=9.5, 9.0, C<sub>3</sub>-H), 7.25-8.12 (15H, m, phenyl). Anal. Calcd. for C<sub>31</sub>H<sub>29</sub>O<sub>10</sub>Cl<sub>3</sub>: C, 55.73; H, 4.34. Found: C, 55.32; H, 4.34.

2,2,2-Trichloroethyl 2,3,4-tri-O-benzoyl-6-O-trityl- $\beta$ -D-glucopyranoside (**31**)

A solution of **14** (444 mg, 0.8 mM) dissolved in anhydrous pyridine (8 ml) was treated with benzoyl chloride (931 μl, 8 mM) at 50°C for 4 h. The solution was worked-up by the standard method. The residual pyridine was removed as the

EtOH azeotrope. The resulting brown oil was purified by P-TLC (EtOAc / n-hexane (1:4)) to give 31 as colorless crystals (604 mg, 87.1% yield). The crystals obtained were recrystallized from EtOH; m.p. 92-94°C;  $[\alpha]_D -5.2^\circ$ ; UV  $\lambda_{\text{max}}^{\text{dioxane}}$  nm( $\epsilon$ ): 275 (3170), 283 (2430); IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 1740 (C=O), 1275 (C-O);  $^1\text{H-NMR}$ : 3.29 (1H, dd,  $J=12.0, 4.0$ ,  $\text{C}_6\text{-H}_a$ ), 3.44 (1H, dd,  $J=12.0, 2.0$ ,  $\text{C}_6\text{-H}_b$ ), 3.85-3.98 (1H, m,  $\text{C}_5\text{-H}$ ), 4.29 (1H, d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.52 (1H, d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 5.15 (1H, d,  $J=7.5$ ,  $\text{C}_1\text{-H}$ ), 5.65 (1H, dd,  $J=9.0, 7.5$ ,  $\text{C}_2\text{-H}$ ), 5.72 (1H, t,  $J=9.0$ ,  $\text{C}_4\text{-H}$ ), 5.84 (1H, t,  $J=9.5$ ,  $\text{C}_3\text{-H}$ ), 7.12-7.98 (30H, m, phenyl). Anal. Calcd. for  $\text{C}_{48}\text{H}_{39}\text{O}_9\text{Cl}_3$ : C, 66.55; H, 4.51. Found: C, 66.54; H, 4.52.

**2,2,2-Trichloroethyl 2,3,6-tri-O-benzoyl- $\beta$ -D-glucopyranoside (32)**

To a solution of 31 (260 mg, 0.3 mM) in  $\text{ClCH}_2\text{CH}_2\text{Cl}$  (3 ml), p-toluenesulfonic acid (206 mg, 1.2 mM) was added. After heating at 60°C for 3 h, the mixture was worked-up by the standard method. The residue was purified by P-TLC (EtOAc / n-hexane (1:4)) to afford 32 (129.5 mg, 69.1% yield), which was crystallized from EtOH / n-hexane; m.p. 192-193°C;  $[\alpha]_D 42.3^\circ$ ; UV  $\lambda_{\text{max}}^{\text{dioxane}}$  nm( $\epsilon$ ): 275 (2760), 282 (2220); IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 3500 (OH), 1725 (C=O), 1280 (C-O);  $^1\text{H-NMR}$ : 3.77-4.02 (1H, m,  $\text{C}_5\text{-H}$ ), 3.96 (1H, t,  $J=8.5$ ,  $\text{C}_4\text{-H}$ ), 4.19 (1H, d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.43 (1H, d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.66 (1H, dd,  $J=12.0, 2.0$ ,  $\text{C}_6\text{-H}_a$ ), 4.83 (1H, dd,  $J=12.0, 4.0$ ,  $\text{C}_6\text{-H}_b$ ), 5.10 (1H, d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ), 5.42-5.63 (2H, m,  $\text{C}_2\text{-H}, \text{C}_3\text{-H}$ ), 7.26-8.12 (15H, m, phenyl). Anal. Calcd.



for  $C_{29}H_{25}O_9Cl_3$ : C, 55.81 ; H, 4.01. Found: C, 55.83 ; H, 3.93.

2,2,2-Trichloroethyl 2,3,6-tri-O-benzoyl-4-O-monochloroacethyl- $\beta$ -D-glucopyranoside (33)

A solution of 32 (50 mg, 0.08 mM) in EtOAc (2 ml) was treated with chloroacetic anhydride (136 mg, 0.8 mM) and pyridine (323  $\mu$ l, 4 mM) at 0°C. After 10 min, the mixture was worked-up by the standard method. The residual pyridine was removed as the EtOH azeotrope. The resulting yellow oil was purified by P-TLC (EtOAc / *n*-hexane (1:4)) to afford colorless crystals (48.1 mg, 81.0% yield), which were recrystallized from EtOH; m.p. 169-170°C;  $[\alpha]_D$  47.1°; UV

$\lambda_{\text{max}}^{\text{dioxane}}$  nm( $\epsilon$ ) : 275 (2840), 283 (2280); IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$  : 1730 (C=O), 1255 (C-O);  $^1\text{H-NMR}$  : 3.93 (1H, s, -OCOCH<sub>2</sub>Cl), 3.94 (1H, s, -OCOCH<sub>2</sub>Cl), 4.11 (1H, ddd, J=9.0, 5.0, 2.5, C<sub>5</sub>-H), 4.19 (1H, d, J=12.0, -OCH<sub>2</sub>CCl<sub>3</sub>), 4.44 (1H, d, J=12.0, -OCH<sub>2</sub>CCl<sub>3</sub>), 4.53 (1H, dd, J=12.5, 4.0, C<sub>6</sub>-H<sub>a</sub>), 4.66 (1H, dd, J=12.5, 2.5, C<sub>6</sub>-H<sub>b</sub>), 5.16 (1H, d, J=8.0, C<sub>1</sub>-H), 5.58 (1H, t, J=9.5, C<sub>4</sub>-H), 5.59 (1H, dd, J=9.0, 7.5, C<sub>2</sub>-H), 5.77 (1H, t, J=9.5, C<sub>3</sub>-H), 7.25-8.11 (15H, m, phenyl). Anal. Calcd. for  $C_{31}H_{26}O_{10}Cl_4$ : C, 53.14 ; H, 3.71. Found: C, 52.64 ; H, 3.60.

2,2,2-Trichloroethyl 2,3,6-tri-O-acetyl-2',3',6'-tri-O-benzyl-4'-O-p-methoxybenzyl- $\beta$ -D-cellobioside (37) and 2,2,2-trichloroethyl 2,3,6-tri-O-acetyl-2',3',6'-tri-O-benzyl-4'-O-p-methoxybenzyl- $\beta$ -D-maltoside (38)

Compound 34 (143 mg, 0.2 mM) and compound 17 (50 mg, 0.1 mM) were dried over  $P_2O_5$  in a vacuum desiccator; and powdered molecular sieves 4A (220 mg) were dried overnight in a  $120^\circ C$  oven before use. A solution of 34 and 17 in anhydrous  $CH_2Cl_2$  (1.5 ml) was added to a stirred suspension of molecular sieves in anhydrous  $CH_2Cl_2$  (0.5 ml) below  $-70^\circ C$ . To the resulting suspension, 0.1 ml of a solution of  $BF_3 \cdot Et_2O$  (2.46  $\mu l$ , 0.02 mM) was added dropwise. The reaction mixture was kept below  $-70^\circ C$  for 2 h, filtered and washed with EtOAc. The combined filtrate and washings were worked-up by the standard method to give a colorless syrup. This syrup was purified by P-TLC (EtOAc / *n*-hexane (1:4) three times and then EtOAc / benzene (5:95) three times) to afford compound 37 (17.3 mg, 16.4% yield) and compound 38 (30.7 mg, 29.0% yield).

Compound 37; m.p.  $86-88^\circ C$ ; UV  $\lambda_{max}^{dioxane}$  nm( $\epsilon$ ): 259 (830), 265(880), 269(900), 275(930), 282(790); IR  $\nu_{max}^{KBr}$   $cm^{-1}$ : 1753 (C=O), 1241 (C-O);  $^1H$ -NMR: 1.99 (3H, s), 2.01 (3H, s), 2.06 (3H, s), 3.77 (3H, s,  $-OC_6H_4OCH_3$ ), 3.81 (1H, t,  $J=9.0$ ,  $C_4-H$ ), 4.09 and 4.35 (2H, two d,  $J=12.0$ ,  $-OCH_2CCl_3$ ), 4.18 (1H, dd,  $J=12.0$ , 4.5,  $C_6-H_a$ ), 4.26 (1H, d,  $J=8.0$ ,  $C_1-H$ ), 4.45 (1H, dd,  $J=12.0$ , 2.0,  $C_6-H_b$ ), 4.77 (1H, d,  $J=8.0$ ,  $C_1-H$ ), 5.04 (1H, dd,  $J=9.0$ , 8.0,  $C_2-H$ ), 5.22 (1H, t,  $J=9.0$ ,  $C_3-H$ ), 6.82 (2H, d,  $J=8.0$ ,  $-OC_6H_4OCH_3$ ), 7.08 (2H, d,  $J=8.0$ ,  $-OC_6H_4OCH_3$ );  $^{13}C$ -NMR: 96.2 ( $-CCl_3$ ), 101.3 (C-1), 102.8 (C-1'), 169.4, 170.1, 170.4 ( $-OAc$ ). Anal. Calcd. for  $C_{49}H_{55}O_{15}Cl_3$ : C, 59.42; H, 5.56. Found: C, 51.02; H, 4.80.

Compound 38: m.p.  $173-174^\circ C$ ; UV  $\lambda_{max}^{dioxane}$  nm( $\epsilon$ ): 260

(1070), 266(1260), 269(1320), 276(1430), 282(1220); IR  $\nu_{\text{max}}^{\text{KBr}}$   
 $\text{cm}^{-1}$ : 1747(C=O), 1235(C-O);  $^1\text{H-NMR}$ : 1.94 (3H, s), 2.01 (3H, s), 2.07 (3H, s), 3.77 (3H, s,  $-\text{OC}_6\text{H}_4\text{OCH}_3$ ), 3.86 (1H, t,  $J=9.0$ ,  $\text{C}_4\text{-H}$ ), 4.10 and 4.37 (2H, two d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.27 (1H, dd,  $J=12.0$ , 4.0,  $\text{C}_6\text{-H}_a$ ), 4.60 (1H, dd,  $J=12.0$ , 2.0,  $\text{C}_6\text{-H}_b$ ), 4.79 (1H, d,  $J=7.5$ ,  $\text{C}_1\text{-H}$ ), 4.80 (1H, d,  $J=3.5$ ,  $\text{C}_1\text{-H}$ ), 5.07 (1H, dd,  $J=9.0$ , 7.5,  $\text{C}_2\text{-H}$ ), 5.32 (1H, t,  $J=9.0$ ,  $\text{C}_3\text{-H}$ ), 6.81 (2H, d,  $J=8.0$ ,  $-\text{OC}_6\text{H}_4\text{OCH}_3$ ), 7.04 (2H, d,  $J=8.0$ ,  $-\text{OC}_6\text{H}_4\text{OCH}_3$ );  $^{13}\text{C-NMR}$ : 96.2 ( $-\text{CCl}_3$ ), 99.1 (C-1'), 101.1 (C-1), 169.4, 169.9, 170.2 ( $-\text{OAc}$ ). Anal. Calcd. for  $\text{C}_{49}\text{H}_{55}\text{O}_{15}\text{Cl}_3$ : C, 59.42; H, 5.56. Found: C, 59.51; H, 5.58.

## 2. EXPERIMENTAL (Chapter 2. Section 2-1)

### 2,2,2-Trichloroethyl 4,6-O-benzylidene- $\beta$ -D-glucopyranoside (42)

To a solution of compound 13 (218 mg, 0.7 mM) and benzaldehyde dimethylacetal<sup>16)</sup> (167  $\mu$ l, 1.1 mM) in dioxane (20 ml), p-toluenesulfonic acid (50 mg, 0.29 mM) was added. The solution was kept at 40°C under 15 mm Hg for 15 min. The solution was worked-up by the standard method to give a colorless oil. The oil was purified by P-TLC (EtOAc / *n*-hexane (1:2)) to afford compound 42, which was crystallized from *iso*-PrOH / *n*-hexane to give colorless crystals (262 mg, 93.6% yield); m.p. 72-74°C;  $[\alpha]_D -41.5^\circ$ ; UV  $\lambda_{\text{max}}^{\text{dioxane}}$  nm ( $\epsilon$ ): 251(160), 257(190), 263(160), 268(70); <sup>1</sup>H-NMR: 2.90 (1H, broad d, J=2.5, C<sub>2</sub>-OH), 3.01 (1H, broad s, C<sub>3</sub>-OH), 3.49 (1H, td, J=9.5, 5.0, C<sub>5</sub>-H), 3.60 (1H, t, J=9.0, C<sub>3</sub>-H), 3.65 (1H, t, J=9.0, C<sub>2</sub>-H), 3.81 (1H, dd, J=10.0, 9.5, C<sub>6</sub>-H<sub>ax</sub>), 3.87 (1H, t, J=9.0, C<sub>4</sub>-H), 4.19 and 4.46 (2H, two d, J=11.0, -OCH<sub>2</sub>CCl<sub>3</sub>), 4.36 (1H, dd, J=10.0, 5.0, C<sub>6</sub>-H<sub>eq</sub>), 4.69 (1H, d, J=7.5, C<sub>1</sub>-H), 5.55 (1H, s, -CHC<sub>6</sub>H<sub>5</sub>), 7.27-7.53 (5H, m, phenyl). Anal. Calcd. for C<sub>15</sub>H<sub>17</sub>O<sub>6</sub>Cl<sub>3</sub>: C, 45.06; H, 4.26. Found: C, 46.24; H, 4.87.

### 2,2,2-Trichloroethyl 2,3-di-O-acetyl-4,6-O-benzylidene- $\beta$ -D-glucopyranoside (43)

Compound 43 was obtained from 42 by the standard acetylation method (Ac<sub>2</sub>O / NaOAc, at 80°C for 1.5 h, a

quantative yield), and was crystallized from EtOH; m.p. 172-173°C;  $[\alpha]_D -77.3^\circ$ ; UV  $\lambda_{\text{max}}^{\text{dioxane}}$  nm ( $\epsilon$ ): 251(160), 257(170), 261(150), 268(95); IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 1760(C=O), 1235 (C-O);  $^1\text{H-NMR}$ : 2.07 (3H, s), 2.08 (3H, s), 3.57 (1H, td,  $J=9.5, 4.5$ ,  $\text{C}_5\text{-H}$ ), 3.75 (1H, t,  $J=9.0$ ,  $\text{C}_6\text{-H}_{\text{ax}}$ ), 3.83 (1H, t,  $J=10.0$ ,  $\text{C}_4\text{-H}$ ), 4.17 and 4.39 (2H, two d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.38 (1H, dd,  $J=10.0, 4.5$ ,  $\text{C}_6\text{-H}_{\text{eq}}$ ), 4.92 (1H, d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ), 5.14 (1H, dd,  $J=9.0, 7.5$ ,  $\text{C}_2\text{-H}$ ), 5.38 (1H, t,  $J=9.5$ ,  $\text{C}_3\text{-H}$ ), 5.52 (1H, s,  $-\text{CHC}_6\text{H}_5$ ), 7.35-7.44 (5H, m, phenyl). Anal. Calcd. for  $\text{C}_{19}\text{H}_{21}\text{O}_8\text{Cl}_3$ : C, 47.16; H, 4.34. Found : C, 47.29; H, 4.49.

**2,2,2-Trichloroethyl 2,3-di-O-acetyl-6-O-benzyl- $\beta$ -D-glucopyranoside (39)**

Compound 43 (193 mg, 0.4 mM) and powdered molecular sieves 4A (200 mg) were dried over  $\text{P}_2\text{O}_5$  in a vacuum desiccator before use. To a stirred suspension of 43 and molecular sieves in anhydrous  $\text{CH}_3\text{CN}$  (3 ml),  $\text{NaCNBH}_3$  (265 mg, 4 mM) was added and then  $\text{TMSCl}$  (0.5 ml, 4 mM) in anhydrous  $\text{CH}_3\text{CN}$  (1 ml) was added dropwise. The reaction mixture was kept at room temperature for 3 h, filtered, and washed with EtOAc. The combined filtrate and washings were worked-up by the standard method to give a colorless oil. The oil was purified by P-TLC (EtOAc / *n*-hexane (1:2)) to afford 39 as a syrup (118 mg, 60.8% yield); m.p. 103-104°C;  $[\alpha]_D -44.8^\circ$ ; IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 3424(OH), 1754 (C=O), 1235(C-O);  $^1\text{H-NMR}$ : 2.06 (3H, s), 2.10 (3H, s), 2.94 (1H, broad d,  $J=3.5$ , -OH), 3.59 (1H, dt,  $J=10.0, 5.0$ ,  $\text{C}_5\text{-H}$ ), 3.75-3.85

(1H, m, C<sub>4</sub>-H), 3.81 (2H, d, J=5.0, C<sub>6</sub>-H<sub>b</sub>), 4.14 (1H, d, J=12.0, -OCH<sub>2</sub>CCl<sub>3</sub>), 4.38 (1H, d, J=12.0, -OCH<sub>2</sub>CCl<sub>3</sub>), 4.58 and 4.66 (2H, two d, J=11.0, benzyl), 4.82 (1H, collapsed d, J=8.0, C<sub>1</sub>-H), 5.01-5.17 (2H, m, C<sub>2</sub>-H, C<sub>3</sub>-H). Anal. Calcd. for C<sub>19</sub>H<sub>23</sub>O<sub>8</sub>Cl<sub>3</sub>: C, 46.96; H, 4.74. Found: C, 46.94; H, 4.70.

2,2,2-Trichloroethyl 2,4,6-tri-O-acetyl-3-O-benzyl-β-D-glucopyranoside (25)

Compound 44 (2.19 g, 5 mM) and powdered molecular sieves 4A (4 g) were dried over P<sub>2</sub>O<sub>5</sub> in a vacuum desiccator before use. To a stirred suspension of compound 44 and molecular sieves in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (15 ml), anhydrous CCl<sub>3</sub>CH<sub>2</sub>OH (2.4 ml, 25 mM) was added, and then BF<sub>3</sub>-Et<sub>2</sub>O (3.1 ml, 25mM) was added dropwise over a period of 30 min at 0°C. The reaction mixture was stirred for an additional 4.5 h at 0°C. The molecular sieves were filtered off and washed with EtOAc. The combined filtrate and washings were worked-up by the standard method to give a colorless oil. The residual CCl<sub>3</sub>CH<sub>2</sub>OH was removed by distillation under reduced pressure. The viscous product was crystallized from EtOH to afford colorless crystals (1.68 g, 63.5% yield), m.p. 98.5-99.5°C; [α]<sub>D</sub> -36.3°; UV λ<sub>max</sub><sup>EtOH</sup> nm (ε): 253(190), 259(230), 265(170); IR ν<sub>max</sub><sup>KBr</sup> cm<sup>-1</sup>: 1747(C=O), 1242(C-O). Anal. Calcd. for C<sub>21</sub>H<sub>25</sub>O<sub>9</sub>Cl<sub>3</sub>: C, 47.77; H, 4.74. Found: C, 48.04; H, 4.78.

2,2,2-Trichloroethyl 3-O-benzyl-β-D-glucopyranoside (45)

To a suspension of compound 25 (527 mg, 1mM) in MeOH (10 ml), 28%-NaOMe in MeOH (1 ml, 5 mM) was added. The reaction mixture was kept for 1 h at 50°C, and was neutralized with 1N-hydrochloric acid using 5% phenolphthalein / ethanol solution as an acid-base indicator, and then concentrated. The product was extracted with EtOAc, washed with a 1N-sodium hydroxide solution and with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated in vacuo to give a colorless syrup (362 mg, 90.2% yield),  $[\alpha]_D -17.9^\circ$  (MeOH); UV  $\lambda_{\text{max}}^{\text{EtOH}}$  nm ( $\epsilon$ ): 253(150), 259 (190), 265(140). Anal. Calcd. for C<sub>15</sub>H<sub>19</sub>O<sub>6</sub>Cl<sub>3</sub>: C, 44.83; H, 4.73. Found: C, 46.45; H, 5.00.

2,2,2-Trichloroethyl  
glucopyranoside (46)

3-O-benzyl-6-O-trityl- $\beta$ -D-

To a solution of compound 45 (922 mg, 2.3 mM) and trityl chloride (1.42 g, 4.6 mM) in anhydrous dioxane (25 ml), Et<sub>3</sub>N (1.6 ml, 11.5 mM) was added dropwise. The solution was stirred at 100°C for 5 h, cooled, and then diluted with EtOAc. The organic layer was washed with a saturated NH<sub>4</sub>Cl solution and with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The product was purified on a silica gel column (Wacogel C-200, 60 g, 18 cm x 3 cm) eluted with EtOAc / n-hexane (1:9) to give a colorless oil (1.23 g, 83.1% yield). The product was crystallized from EtOH; m.p. 74-75°C;  $[\alpha]_D -31.3^\circ$ ; UV  $\lambda_{\text{max}}^{\text{dioxane}}$  nm ( $\epsilon$ ): 254(920), 259(970). Anal. Calcd. for C<sub>34</sub>H<sub>33</sub>O<sub>6</sub>Cl<sub>3</sub>: C, 63.40; H, 5.13. Found: C, 63.34; H, 5.12.

2,2,2-Trichloroethyl 2,4-di-O-acetyl-3-O-benzyl-6-O-trityl-

$\beta$ -D-glucopyranoside (47)

Compound 47 was obtained from 46 by the standard acetylation method ( $\text{Ac}_2\text{O}$  /  $\text{NaOAc}$  at  $100^\circ\text{C}$  for 1 h, a quantitative yield), and was crystallized from  $\text{EtOH}$  / n-hexane; m.p.  $144\text{--}145^\circ\text{C}$ ;  $[\alpha]_{\text{D}} -5.3^\circ$ ; UV  $\lambda_{\text{max}}^{\text{dioxane}}$  nm ( $\epsilon$ ): 254(990), 259(1040); IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 1754( $\text{C}=\text{O}$ ), 1222( $\text{C}-\text{O}$ );  $^1\text{H-NMR}$ : 1.70 (3H, s), 2.05 (3H, s), 3.12 (1H, dd,  $J=10.5$ , 5.0,  $\text{C}_6\text{-H}_a$ ), 3.21 (1H, dd,  $J=10.5$ , 2.5,  $\text{C}_6\text{-H}_b$ ), 3.45–3.56 (1H, m,  $\text{C}_5\text{-H}$ ), 3.68 (1H, t,  $J=9.5$ ,  $\text{C}_3\text{-H}$ ), 4.22 and 4.46 (2H, two d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.56 and 4.62 (2H, two d,  $J=12.0$ , benzyl), 4.77 (1H, d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ), 5.20 (1H, t,  $J=9.3$ ,  $\text{C}_4\text{-H}$ ), 5.21 (1H, dd,  $J=9.5$ , 8.0,  $\text{C}_2\text{-H}$ ), 7.19–7.49 (20H, m, phenyl). Anal. Calcd. for  $\text{C}_{38}\text{H}_{37}\text{O}_8\text{Cl}_3$ : C, 62.68; H, 5.09. Found: C, 63.57; H, 5.39.

2,2,2-Trichloroethyl 2,6-di-O-acetyl-3-O-benzyl- $\beta$ -D-glucopyranoside (40)

To a solution of compound 47 (949 mg, 1.27 mM) in  $\text{CH}_2\text{Cl}_2$  (30 ml), *p*-toluenesulfonic acid (437 mg, 2.54 mM) was added. The solution was stirred at room temperature for 3 h, and then worked-up by the standard method to give a yellow oil. The oil was purified by P-TLC ( $\text{EtOAc}$  / n-hexane (1:4)) to afford compound 40 (504.6 mg, 78.6% yield), which was crystallized from  $\text{EtOH}$  / n-hexane; m.p.  $110\text{--}111^\circ\text{C}$ ;  $[\alpha]_{\text{D}} -43.0^\circ$ ; UV  $\lambda_{\text{max}}^{\text{EtOH}}$  nm ( $\epsilon$ ): 252(140), 258(170), 265(130); IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 3424(OH), 1747( $\text{C}=\text{O}$ ), 1248( $\text{C}-\text{O}$ );  $^1\text{H-NMR}$ : 2.05 (3H, s), 2.12 (3H, s), 3.43–3.55 (1H, m,  $\text{C}_5\text{-H}$ ), 3.53–



3.65 (2H, m, C<sub>3</sub>-H, C<sub>4</sub>-H), 4.11 and 4.38 (2H, two d, J=12.0, -OCH<sub>2</sub>CCl<sub>3</sub>), 4.30 (1H, dd, J=12.0, 2.0, C<sub>6</sub>-H<sub>a</sub>), 4.47 (1H, d, J=12.0, 4.0, C<sub>6</sub>-H<sub>b</sub>), 4.71 and 4.78 (2H, two d, J=11.5, benzyl), 4.72 (1H, d, J=8.0, C<sub>1</sub>-H), 5.07 (1H, collapsed dd, J=9.5, 8.0, C<sub>2</sub>-H), 7.28-7.48 (5H, m, phenyl). Anal. Calcd. for C<sub>19</sub>H<sub>23</sub>O<sub>8</sub>Cl<sub>3</sub>: C, 46.96; H, 4.74. Found: C, 47.14; H, 4.80.

**2,2,2-Trichloroethyl 3-O-benzyl-4,6-O-benzylidene-β-D-glucopyranoside (48)**

Compound 48 was obtained from compound 45 in the same way that compound 42 was obtained from compound 13, (PhCH(OMe)<sub>2</sub> / p-TsOH / dioxane at 40°C under 15 mmHg for 20 min, 89.3% yield); m.p. 103-104°C; [α]<sub>D</sub> -37.3°; UV λ<sub>dioxane max</sub> nm (ε): 252(230), 258(280), 262(260); <sup>1</sup>H-NMR: 2.61 (1H, broad s, -OH), 3.48 (1H, m, C<sub>5</sub>-H), 3.67-3.75 (3H, m, C<sub>2</sub>-H, C<sub>3</sub>-H, C<sub>4</sub>-H), 3.81 (1H, t, J=10.0, C<sub>6</sub>-H<sub>ax</sub>), 4.18 and 4.44 (2H, two d, J=11.5, -OCH<sub>2</sub>CCl<sub>3</sub>), 4.36 (1H, dd, J=10.0, 5.0, C<sub>6</sub>-H<sub>eq</sub>), 4.67 (1H, collapsed d, J=7.5, C<sub>1</sub>-H), 4.82 and 4.92 (2H, two d, J=11.5, benzyl), 5.58 (1H, s, -CHC<sub>6</sub>H<sub>5</sub>), 7.25-7.53 (10H, m, phenyl). Anal. Calcd. for C<sub>22</sub>H<sub>23</sub>O<sub>6</sub>Cl<sub>3</sub>: C, 53.93; H, 4.70. Found: C, 53.81; H, 4.72.

**2,2,2-Trichloroethyl 2-O-acetyl-3-O-benzyl-4,6-O-benzylidene-β-D-glucopyranoside (49)**

Compound 49 was prepared from 48 by the standard acetylation method (Ac<sub>2</sub>O / NaOAc at 80°C for 4 h, a quantitative yield), and was crystallized from EtOH; m.p.

132-133°C;  $[\alpha]_D -21.7^\circ$ ; UV  $\lambda_{\text{max}}^{\text{dioxane}}$  nm ( $\epsilon$ ): 252(130), 258(180), 263(140); IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 1747(C=O), 1242(C-O);  $^1\text{H-NMR}$ : 2.02 (3H, s), 3.47 (1H, t,  $J=9.0$ ,  $\text{C}_5\text{-H}$ ), 3.75 (1H, t,  $J=8.5$ ,  $\text{C}_3\text{-H}$ ), 3.81 (1H, t,  $J=9.0$ ,  $\text{C}_4\text{-H}$ ), 3.83 (1H, t,  $J=10.0$ ,  $\text{C}_6\text{-H}_{\text{ax}}$ ), 4.12 and 4.36 (2H, two d,  $J=12.0$ ,  $\text{OCH}_2\text{CCl}_3$ ), 4.36 (1H, dd,  $J=10.0$ , 5.0,  $\text{C}_6\text{-H}_a$ ), 4.68 and 4.88 (2H, two d,  $J=12.0$ , benzyl), 4.78 (1H, d,  $J=7.5$ ,  $\text{C}_1\text{-H}$ ), 5.12 (1H, collapsed t,  $J=8.5$ ,  $\text{C}_2\text{-H}$ ), 5.59 (1H, s,  $-\text{CHC}_6\text{H}_5$ ), 7.29-7.47 (10H, m, phenyl). Anal. Calcd. for  $\text{C}_{24}\text{H}_{25}\text{O}_7\text{Cl}_3$ : C, 54.19; H, 4.70. Found: C, 54.28; H, 4.68.

**2,2,2-Trichloroethyl 2-O-acetyl-3,6-di-O-benzyl- $\beta$ -D-glucopyranoside (41)**

Compound 41 was obtained as a syrup from 49 in the same way that compound 39 was obtained from compound 43, ( $\text{NaCNBH}_3$  /  $\text{TMSCl}$  / anhydrous  $\text{CH}_3\text{CN}$  at r.t. for 15 h, about 60% yield);  $[\alpha]_D -23.2^\circ$ ; UV  $\lambda_{\text{max}}^{\text{dioxane}}$  nm ( $\epsilon$ ): 252(340), 258(420), 263(320), 266(300), 268(210); IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 3488(OH), 1747(C=O), 1241(C-O);  $^1\text{H-NMR}$ : 2.04 (3H, s), 2.68 (1H, broad s, -OH), 3.49 (1H, dt,  $J=10.0$ , 5.0,  $\text{C}_5\text{-H}$ ), 3.55 (1H, t,  $J=9.0$ ,  $\text{C}_3\text{-H}$ ), 3.70 (1H, broad t,  $J=10.0$ ,  $\text{C}_4\text{-H}$ ), 3.76 (2H, d,  $J=5.0$ ,  $\text{C}_6\text{-H}$ ), 4.12 and 4.37 (2H, two d,  $J=11.5$ ,  $\text{OCH}_2\text{CCl}_3$ ), 4.56 and 4.63 (2H, two d,  $J=12.0$ , benzyl), 4.71 (1H, d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ), 4.71 and 4.77 (2H, two d,  $J=11.5$ , benzyl), 5.08 (1H, dd,  $J=9.5$ , 8.0,  $\text{C}_2\text{-H}$ ). Anal. Calcd. for  $\text{C}_{24}\text{H}_{27}\text{O}_7\text{Cl}_3$ : C, 53.98; H, 5.06. Found: C, 54.13; H, 5.10.

### Glycosylation between imidate 34 and aglycons 39-41

Imidate 34 (143 mg, 0.2mM) and aglycon (0.1 mM) were dried over  $P_2O_5$  in a vacuum desiccator; and powdered molecular sieves 4A (220 mg) were dried overnight in a  $120^\circ C$  oven before use. A solution of imidate and aglycon in anhydrous  $CH_2Cl_2$  (1.5 ml) was added to a stirred suspension of molecular sieves in anhydrous  $CH_2Cl_2$  (0.5 ml) cooled below  $-70^\circ C$ . To this reaction mixture,  $BF_3 \cdot Et_2O$  (2.46  $\mu l$ , 0.02 mM) in anhydrous  $CH_2Cl_2$  (0.1 ml) was added dropwise over a period of 10 min below  $-70^\circ C$ . The reaction mixture was then kept below  $-70^\circ C$  for the prescribed time (Table 1), and then filtered. The residual molecular sieves were washed with EtOAc. The combined filtrate and washings were worked-up by the standard method to give a yellow oil. The oil was purified by P-TLC developed with the following solvent combinations (v/v) to afford the corresponding glycoside: aglycon 39, EtOAc / *n*-hexane (1:4) twice and then EtOAc / benzene (5:95) five times; aglycon 40, EtOAc / *n*-hexane (1:2) twice; aglycon 41, EtOAc / *n*-hexane (1:4) twice.

**Compound 51** ( $\beta$ -anomer, 13.0% yield); m.p.  $85-87^\circ C$ ;  
 UV  $\lambda_{max}^{dioxane}$  nm ( $\epsilon$ ): 254(1140), 259(1450), 263(1540), 276(1540); IR  $\nu_{max}^{KBr}$   $cm^{-1}$ : 1760(C=O), 1248(C-O);  $^1H$ -NMR: 1.99 (3H, s), 2.06 (3H, s), 3.78 (3H, s,  $-OC_6H_4OCH_3$ ), 3.97 (1H, t,  $J=9.5$ ,  $C_4-H$ ), 4.12 and 4.38 (2H, two d,  $J=12.0$ ,  $-OCH_2CCl_3$ ), 4.31 (1H, d,  $J=8.0$ ,  $C_1-H$ ), 4.78 (1H, d,  $J=7.5$ ,  $C_1-H$ ), 5.08 (1H, dd,  $J=9.5$ ,  $7.5$ ,  $C_2-H$ ), 5.22 (1H, t,  $J=9.5$ ,  $C_3-H$ ), 6.80 (2H, d,  $J=8.0$ ,  $-OC_6H_4OCH_3$ ), 7.10 (2H, d,  $J=8.0$ ,  $-OC_6H_4OCH_3$ ). Anal. Calcd. for  $C_{54}H_{59}O_{14}Cl_3$ : C, 62.46; H,

5.69. Found: C, 62.38; H, 5.58.

**Glycoside 52** ( $\alpha$ -anomer, 33.0% yield); UV  $\lambda_{\text{max}}^{\text{dioxane}}$  nm ( $\epsilon$ ): 254(1890), 259(2110), 269(2110); IR  $\nu_{\text{max}}^{\text{kBr}}$   $\text{cm}^{-1}$ : 1760(C=O), 1248(C-O);  $^1\text{H-NMR}$ : 1.90 (3H, s), 2.06 (3H, s), 3.79 (3H, s,  $-\text{OC}_6\text{H}_4\text{OCH}_3$ ), 3.96 (1H, dd,  $J=11.0, 4.0$ ,  $\text{C}_6\text{-H}_a$ ), 4.08 (1H, t,  $J=9.0$ ,  $\text{C}_4\text{-H}$ ), 4.13 and 4.40 (2H, two d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.78 (1H, d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ), 5.01 (1H, d,  $J=4.0$ ,  $\text{C}_1\text{-H}$ ), 5.08 (1H, dd,  $J=9.0, 8.0$ ,  $\text{C}_2\text{-H}$ ), 5.35 (1H, t,  $J=9.0$ ,  $\text{C}_3\text{-H}$ ), 6.80 (2H, d,  $J=8.0$ ,  $-\text{OC}_6\text{H}_4\text{OCH}_3$ ), 7.02 (2H, d,  $J=8.0$ ,  $-\text{OC}_6\text{H}_4\text{OCH}_3$ ). Anal. Calcd. for  $\text{C}_{54}\text{H}_{59}\text{O}_{14}\text{Cl}_3$ : C, 62.46; H, 5.69. Found: C, 61.97; H, 6.41.

**Glycoside 53** ( $\beta$ -anomer, 94.2% yield); m.p.  $105^\circ\text{C}$ ;  $[\alpha]_D^{20}$ : 2.9 $^\circ$ ; UV  $\lambda_{\text{max}}^{\text{dioxane}}$  nm ( $\epsilon$ ): 260(1510), 266(1710), 269(1740), 275(1790), 282(1510); IR  $\nu_{\text{max}}^{\text{kBr}}$   $\text{cm}^{-1}$ : 1747(C=O), 1235(C-O);  $^1\text{H-NMR}$ : 1.93 (3H, s), 2.01 (3H, s), 3.77 (3H, s,  $-\text{OC}_6\text{H}_4\text{OCH}_3$ ), 3.93 (1H, t,  $J=9.0$ ,  $\text{C}_4\text{-H}$ ), 4.05 and 4.32 (2H, two d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.23 (1H, dd,  $J=12.0, 4.5$ ,  $\text{C}_6\text{-H}_a$ ), 4.47 (1H, d,  $J=10.0$ ,  $\text{C}_1\text{-H}$ ), 4.49 (1H, dd,  $J=12.0, 2.5$ ,  $\text{C}_6\text{-H}_b$ ), 4.67 (1H, d,  $J=9.0$ ,  $\text{C}_1\text{-H}$ ), 5.04 (1H, dd,  $J=9.0, 7.5$ ,  $\text{C}_2\text{-H}$ ), 6.83 (2H, d,  $J=8.0$ ,  $-\text{OC}_6\text{H}_4\text{OCH}_3$ ), 7.12 (2H, d,  $J=8.0$ ,  $-\text{OC}_6\text{H}_4\text{OCH}_3$ );  $^{13}\text{C-NMR}$ : 96.5 ( $-\text{CCl}_3$ ), 101.4(C-1), 102.8(C-1'), 169.3, 170.3 ( $-\text{OAc}$ ). Anal. Calcd. for  $\text{C}_{54}\text{H}_{59}\text{O}_{14}\text{Cl}_3$ : C, 62.46; H, 5.69. Found: C, 62.70; H, 5.70.

**Glycoside 54** ( $\beta$ -anomer, 96.0% yield);  $[\alpha]_D^{20}$ : 3.9 $^\circ$ ; UV  $\lambda_{\text{max}}^{\text{dioxane}}$  nm ( $\epsilon$ ): 254(970), 260(1240), 265(1310), 278(1240); IR  $\nu_{\text{max}}^{\text{kBr}}$   $\text{cm}^{-1}$ : 1754(C=O), 1235(C-O);  $^1\text{H-NMR}$ : 1.93 (3H, s), 3.78 (3H, s,  $-\text{OC}_6\text{H}_4\text{OCH}_3$ ), 4.09 and 4.35 (2H, two d,

$J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ),  $4.47$  (1H, d,  $J=9.0$ ,  $\text{C}_1\text{-H}$ ),  $4.67$  (1H, d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ),  $5.06$  (1H, dd,  $J=9.0$ ,  $8.0$ ,  $\text{C}_2\text{-H}$ ),  $6.79$  (2H, d,  $J=8.0$ ,  $-\text{OC}_6\text{H}_4\text{OCH}_3$ ),  $7.07$  (2H, d,  $J=8.0$ ,  $-\text{OC}_6\text{H}_4\text{OCH}_3$ );  $^{13}\text{C}$ -NMR:  $96.6$  ( $-\text{CCl}_3$ ),  $101.6$  ( $\text{C-1}$ ),  $102.6$  ( $\text{C-1'}$ ),  $169.3$  ( $-\text{OAc}$ ).  
 Anal. Calcd. for  $\text{C}_{59}\text{H}_{63}\text{O}_{13}\text{Cl}_3$ : C, 65.22; H, 5.80. Found. C, 64.92; H, 5.61.

### 3. EXPERIMENTAL (Chapter 2. Section 2-2)

2,2,2-Trichloroethyl    2,4,6-tri-O-acetyl-3-O-propargyl- $\beta$ -D-glucopyranoside (59)

Compound 59 was prepared from compound 58 in the same way that compound 25 was prepared from compound 44, ( $\text{CCl}_3\text{CH}_2\text{OH}$  /  $\text{BF}_3\text{-Et}_2\text{O}$  / anhydrous  $\text{CH}_2\text{Cl}_2$ , at  $0^\circ\text{C}$  for 15 h, 57.9% yield).

2,2,2-Trichloroethyl 3-O-propargyl- $\beta$ -D-glucopyranoside (60)

Compound 60 was obtained from compound 59 in the same way that compound 45 was obtained from compound 25, (28%-NaOMe in MeOH / MeOH, at  $50^\circ\text{C}$  for 3 h, a quantitative yield).

2,2,2-Trichloroethyl                    3-O-propargyl-6-O-trityl- $\beta$ -D-glucopyranoside (61)

Compound 61 was synthesized from compound 60 in the same way that compound 46 was synthesized from compound 45, ( $\text{TrCl}$  /  $\text{Et}_3\text{N}$  / anhydrous dioxane, at  $100^\circ\text{C}$  for 3 h, 73.0% yield).

2,2,2-Trichloroethyl                    2,4-di-O-acetyl-3-O-propargyl-6-O-trityl- $\beta$ -D-glucopyranoside (62)

Compound 62 was obtained from compound 61 by the standard acetylation method, ( $\text{Ac}_2\text{O}$  /  $\text{AcONa}$ , at  $100^\circ\text{C}$  for 1 h, a quantitative yield).

2,2,2-Trichloroethyl                    2,6-di-O-acetyl-3-O-propargyl- $\beta$ -D-

glucopyranoside (57)

To a solution of compound 62 (1427 mg, 2.1 mM) in  $\text{CH}_2\text{Cl}_2$  (20 ml), p-toluenesulfonic acid (730 mg, 4.2mM) was added. The solution was stirred at room temperature for 2.5 h, and then worked-up by the standard method to give a colorless oil. The oil was purified by P-TLC (EtOAc / n-hexane (1:2)) to afford compound 57 (764 mg, 83.2% yield), which was crysallized from EtOH / n-hexane; m.p. 126-127°C;  $^1\text{H}$ -NMR: 2.15 (3H, s), 2.16 (3H, s), 2.51 (1H, t,  $J=2.5$ ,  $-\text{C}=\text{CH}$ ), 3.04 (1H, d,  $J=2.0$ ,  $-\text{OH}$ ), 3.46-3.72 (3H, m,  $\text{C}_3\text{-H}$ ,  $\text{C}_4\text{-H}$ ,  $\text{C}_5\text{-H}$ ), 4.13 and 4.40 (2H, two d.  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.31 and 4.47 (2H, two dd,  $J=17.0$ , 2.5,  $-\text{CH}_2\text{C}=\text{C}$ ), 4.33 (1H, dd,  $J=12.0$ , 2.0,  $\text{C}_6\text{-H}_a$ ), 4.48 (1H, dd,  $J=12.0$ , 3.5,  $\text{C}_6\text{-H}_b$ ), 4.75 (1H, d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ), 5.02 (1H, dd,  $J=9.0$ , 8.0,  $\text{C}_2\text{-H}$ ). Anal. Calcd. for  $\text{C}_{15}\text{H}_{19}\text{O}_8\text{Cl}_3$ : C, 41.52; H, 4.38. Found: C, 41.45, H, 4.50.

2,2,2-Trichloroethyl  
glucopyranoside (56)

2,6-di-O-acetyl-3-O-allyl- $\beta$ -D-

A solution of compound 57 (15 mg, 0.03 mM) in MeOH (0.7 ml) was treated with 5% Pd/BaSO<sub>4</sub> (1 mg) under H<sub>2</sub> at 0°C for 30 min. The reaction mixture was filtered, and evaporated in vacuo to give a oil. The oil was purified by P-TLC (EtOAc / n-hexane (1:2) twice), and then was crystallized from EtOH / n-hexane to give compound 56 (4.6 mg, 30.5% yield) as colorless crystals; m.p. 123-124°C;  $^1\text{H}$ -NMR: 2.11 (3H, s), 2.13 (3H, s), 2.90 (1H, d,  $J=4.0$ ,  $-\text{OH}$ ), 3.36-3.63 (3H, m,

$C_3$ -H,  $C_4$ -H,  $C_5$ -H), 4.11 and 4.38 (2H, two d,  $J=12.0$ ,  $-OCH_2CCl_3$ ), 4.21 and 4.22 (2H, two d,  $J=5.5$ ,  $-CH_2-C=C$ ), 4.32 (1H, dd,  $J=12.0$ , 2.0,  $C_6-H_a$ ), 4.47 (1H, dd,  $J=12.0$ , 2.0,  $C_6-H_b$ ), 4.72 (1H, d,  $J=8.0$ ,  $C_1$ -H), 5.02 (1H, dd,  $J=9.0$ , 8.0,  $C_2$ -H), 5.18 (1H, dd,  $J=10.5$ , 1.5,  $-C-C=CH_2$  cis), 5.26 (1H, dd,  $J=17.5$ , 1.5,  $-C-C=CH_2$  trans), 5.87 (1H, dddd,  $J=17.5$ , 10.5, 5.5, 5.5,  $-C-CH=C$ ). Anal. Calcd. for  $C_{15}H_{21}O_8Cl_3$ : C, 41.33; H, 4.82. Found: C, 41.29; H, 4.81.

**2,2,2-Trichloroethyl 2,6-di-O-acetyl-3-O-propyl- $\beta$ -D-glucopyranoside (55)**

A solution of compound 57 (94 mg, 0.2mM) in MeOH (2 ml) was treated with Pd-C (63 mg) under  $H_2$  at  $0^\circ C$  for 30 min. The reaction mixture was filtered and concentrated to afford a oil. The oil was crystallized from EtOH / n-hexane (56.2mg, 59.1% yield); m.p. 118-119 $^\circ C$ ;  $^1H$ -NMR: 0.91 (3H, t,  $J=7.2$ ,  $-C-C-CH_3$ ), 1.58 (2H, dddd,  $J=7.2$ , 7.2, 7.2, 7.2,  $-C-CH_2-C$ ), 2.12 (3H, s), 2.14 (3H, s), 2.85 (1H, broad d,  $J=3.0$ ,  $-OH$ ), 3.39 (1H, t,  $J=9.5$ ,  $C_3$ -H), 3.46-3.73 (4H, m,  $C_4$ -H,  $C_5$ -H,  $-CH_2-C-C$ ), 4.12 and 4.39 (2H, two d,  $J=12.0$ ,  $-OCH_2CCl_3$ ), 4.33 (1H, dd,  $J=12.0$ , 2.0,  $C_6-H_a$ ), 4.45 (1H, dd,  $J=12.0$ , 4.0,  $C_6-H_b$ ), 4.72 (1H, d,  $J=8.5$ ,  $C_1$ -H), 5.00 (1H, dd,  $J=9.5$ , 8.5,  $C_2$ -H). Anal. Calcd. for  $C_{15}H_{23}O_8Cl_3$ : C, 41.14; H, 5.26. Found: C, 41.07; H, 5.27.

**2,2,2-Trichloroethyl 4,6-O-benzylidene-2,3-di-O-carbamoyl- $\beta$ -D-glucopyranoside (70)**

To a solution of compound 42 (2.36 g, 5.9mM) in pyridine



(15 ml), PhNCO (3.2 ml, 29.5 mM) was added. The solution was kept at 100°C for 5 h. The reaction did not complete, and then PhNCO (1.6 ml, 14.75 mM) was added again. The reaction mixture was kept at 100°C for another 1 h. The excess PhNCO was decomposed by the addition of MeOH (2 ml) at 100°C for 30 min. The reaction mixture was diluted with EtOAc, washed with a 1N-hydrochloric acid solution and with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and evaporated in vacuo. The product was extracted with CH<sub>2</sub>Cl<sub>2</sub>, and then filtered. The filtrate was concentrated to give colorless crystals (2.72 g, 72.2% yield).

**2,2,2-Trichloroethyl 4,6-O-benzylidene-2,3-di-O-pivaloyl-β-D-glucopyranoside (71)**

To a solution of compound 42 (582 mg, 1.45 mM) in pyridine (7 ml), (CH<sub>3</sub>)<sub>3</sub>CCOCl (1.8 ml, 14.5 mM) was added. The solution was kept at 50°C for 15 h, and worked-up in the same way as that for compound 70 to give a oil. The product was crystallized from EtOH to afford colorless crystals (596 mg, 72.1% yield).

**2,2,2-Trichloroethyl 2,3-di-O-carbamoyl-β-D-glucopyranoside (72)**

To a solution of compound 70 (2.7 g, 4.2mM) in 20%-MeOH / CH<sub>2</sub>Cl<sub>2</sub> (20 ml), BF<sub>3</sub>-Et<sub>2</sub>O (2.58 mM, 21mM) was added at 0°C. The solution was stirred at room temperature for 1.5 h, and worked-up by the standard method to afford colorless

crystals which were recrystallized from EtOH / n-hexane (1.7g, 73.0% yield).

2,2,2-Trichloroethyl 2,3-di-O-pivaloyl- $\beta$ -D-glucopyranoside (73)

Compound 73 was obtained from compound 71 in the same way that compound 72 was obtained from compound 70, ( $\text{BF}_3\text{-Et}_2\text{O}$  / 20%-MeOH /  $\text{CH}_2\text{Cl}_2$ , at r.t. for 3 h, 61.5% yield).

2,2,2-Trichloroethyl 2,3,6-tri-O-carbamoyl- $\beta$ -D-glucopyranoside (66)

To a solution of compound 72 (60 mg, 0.11 mM) in anhydrous pyridine (2 ml),  $\text{PhNCO}$  (96  $\mu\text{l}$ , 0.88 mM) was added. The solution was kept at 50°C for 30 min. The excess  $\text{PhNCO}$  was decomposed by the addition of MeOH (1 ml) at 50°C for 30 min. The solution was extracted with EtOAc, washed with a 1N-hydrochloric acid solution and with brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated to give colorless crystals. The product was purified on silica gel column (Wacogel C-200, 24g, 2.5cm x 9.0cm) eluted with EtOAc/n-hexane (1:2) to give compound 66 as crystals (43 mg, 58.9% yield); m.p. 199-200°C;  $^1\text{H-NMR}$ : 3.20-3.32 (1H, m,  $\text{C}_5\text{-H}$ ), 3.30-3.60 (1H, broad s, -OH), 3.60 (1H, t,  $J=7.5$ ,  $\text{C}_4\text{-H}$ ), 4.10 and 4.33 (2H, two d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.36 (1H, broad d,  $J=11.0$ ,  $\text{C}_6\text{-H}_a$ ), 4.43 (1H, broad d,  $J=11.0$ ,  $\text{C}_6\text{-H}_b$ ), 4.81 (1H, d,  $J=7.5$ ,  $\text{C}_1\text{-H}$ ), 4.86 (1H, dd,  $J=8.0$ , 7.5,  $\text{C}_2\text{-H}$ ), 4.96 (1H, dd,  $J=8.0$ , 7.5,  $\text{C}_3\text{-H}$ ), 6.94-7.13 (3H, m,  $-\text{CONHPh}$ ), 7.16-7.31 (15H, m, phenyl). Anal. Calcd. for  $\text{C}_{29}\text{H}_{28}\text{O}_9\text{Cl}_3\text{N}_3$ : C, 52.06; H, 4.19.

Found: C, 51.79; H, 4.17.

2,2,2-Trichloroethyl                      6-O-acetyl-2,3-di-O-carbamoyl- $\beta$ -D-  
glucopyranoside (67)

To a solution of compound 72 (121 mg, 0.22 mM) in EtOAc (3 ml),  $\text{CH}_3\text{COCl}$  (63  $\mu\text{l}$ , 0.88 mM) and 2,6-lutidine (128  $\mu\text{l}$ , 1.1 mM) were added. The solution was stirred at  $0^\circ\text{C}$  for 4 h, and then worked up in the same way as that for 66 to give a colorless oil. The oil was purified by P-TLC ( $\text{MeOH}/\text{CH}_2\text{Cl}_2$  (5:95)) to afford compound 67 (84 mg, 64.5% yield), which was crystallized from EtOH/n-hexane; m.p.  $157\text{--}159^\circ\text{C}$ ;  $^1\text{H-NMR}$ : 2.14 (3H, s), 3.55–3.80 (2H, m,  $\text{C}_4\text{-H}$ ,  $\text{C}_5\text{-H}$ ), 4.18 and 4.42 (2H, two d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.41 (1H, broad s,  $\text{C}_6\text{-H}_a$ ), 4.43 (1H, broad s,  $\text{C}_6\text{-H}_b$ ), 4.88 (1H, d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ), 4.95–5.15 (2H, m,  $\text{C}_2\text{-H}$ ,  $\text{C}_3\text{-H}$ ), 7.00–7.12 (2H, m,  $-\text{CONHPh}$ ), 7.14–7.40 (10H, m, phenyl). Anal. Calcd. for  $\text{C}_{24}\text{H}_{25}\text{O}_9\text{Cl}_3\text{N}_2$ : C, 48.69; H, 4.23. Found: C, 49.38; H, 4.42.

2,2,2-Trichloroethyl                      2,3,6-tri-O-pivaloyl- $\beta$ -D-  
glucopyranoside (68)

To a solution of compound 73 (192 mg, 0.4 mM) in  $\text{CHCl}_3$  (7 ml),  $(\text{CH}_3)_3\text{CCOCl}$  (197  $\mu\text{l}$ , 1.6 mM) and pyridine (162  $\mu\text{l}$ , 2.0 mM) were added. The solution was kept at  $50^\circ\text{C}$  for 30hr, and then worked up in the same way as that for 66 to afford a colorless oil. The oil was crystallized from EtOH / n-hexane to give compound 68 (145 mg, 64.3% yield); m.p.  $194\text{--}195^\circ\text{C}$ ;  $^1\text{H-NMR}$ : 1.20 (9H, s), 1.22 (9H, s), 1.26 (9H, s), 3.15 (1H, d,  $J=5.0$ ,  $-\text{OH}$ ), 3.45–3.66 (2H, m,  $\text{C}_4\text{-H}$ ,  $\text{C}_5\text{-H}$ ),

4.14 and 4.32 (2H, two d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.37-4.44 (2H, broad,  $\text{C}_6\text{-H}_a$ ,  $\text{C}_6\text{-H}_b$ ), 4.87 (1H, collapsed d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ), 5.00-5.06 (2H, m,  $\text{C}_2\text{-H}$ ,  $\text{C}_3\text{-H}$ ). Anal. Calcd. for  $\text{C}_{23}\text{H}_{37}\text{O}_9\text{Cl}_3$ : C, 48.98; H, 6.57. Found: C, 48.80; H, 6.67.

2,2,2-Trichloroethyl 2,3-di-O-monochloroacetyl-6-O-trityl- $\beta$ -D-glucopyranoside (74)

To a solution of compound 14 (101 mg, 0.18 mM) in EtOAc (4 ml),  $(\text{ClCH}_2\text{CO})_2\text{O}$  (468 mg, 2.74 mM) and pyridine (0.74 ml, 9.13 mM) were added at  $0^\circ\text{C}$ . The reaction mixture was stirred at  $0^\circ\text{C}$  for 20 min, and partitioned between EtOAc and brine. The EtOAc layer was dried over  $\text{Na}_2\text{SO}_4$  and evaporated in vacuo to give an yellow oil which was purified by P-TLC (EtOAc / n-hexane (1:2)) to afford a colorless oil (107 mg, 75.0% yield).

2,2,2-Trichloroethyl 2,3,6-tri-O-monochloroacetyl- $\beta$ -D-glucopyranoside (69)

To a solution of compound 74 (97 mg, 0.12mM) in  $\text{CH}_2\text{Cl}_2$  (3.5 ml), *p*-toluenesulfonic acid (43 mg, 0.25 mM) was added. The reaction mixture was kept at room temperature for 2.5 h, and worked-up by the standard method. The product was purified by P-TLC (EtOAc / n-hexane (1:2)) to afford a colorless oil (47.1 mg, 70.1% yield); m.p.  $125\text{-}126^\circ\text{C}$ ;  $^1\text{H}$ -NMR: 3.60 -3.71 (1H, m,  $\text{C}_5\text{-H}$ ), 3.73 (1H, dd,  $J=9.5$ ,  $8.0$ ,  $\text{C}_4\text{-H}$ ), 4.06 (2H, s,  $-\text{COCH}_2\text{Cl}$ ), 4.08 and 4.13 (2H, two d,  $J=9.0$ ,  $-\text{COCH}_2\text{Cl}$ ), 4.15 and 4.39 (2H, two d,  $J=12.0$ ,  $-\text{CH}_2\text{CCl}_3$ ), 4.16

(2H, s,  $-\text{COCH}_2\text{Cl}$ ), 4.50-4.56 (2H, m,  $\text{C}_6\text{-H}_a$ ,  $\text{C}_6\text{-H}_b$ ), 4.91 (1H, d,  $J=7.5$ ,  $\text{C}_1\text{-H}$ ), 5.11 (1H, dd,  $J=9.5$ ,  $7.5$ ,  $\text{C}_2\text{-H}$ ), 5.22 (1H, dd,  $J=9.5$ ,  $8.0$ ,  $\text{C}_3\text{-H}$ ). Anal. Calcd. for  $\text{C}_{14}\text{H}_{16}\text{O}_9\text{Cl}_6$ : C, 31.05; H, 2.96. Found: C, 31.20; H, 2.92.

#### Glycosylation between imidate 34 and aglycons 55-57 and 66-69

Imidate 34 (2.0eq) and aglycon (1.0eq) were dried over  $\text{P}_2\text{O}_5$  in a vacuum desiccator; powdered molecular sieves 4A were dried overnight in a  $120^\circ\text{C}$  oven before use. A solution of imidate and aglycon in anhydrous  $\text{CH}_2\text{Cl}_2$  was added to a stirred suspension of molecular sieves in anhydrous  $\text{CH}_2\text{Cl}_2$  below  $-70^\circ\text{C}$ . To this reaction mixture,  $\text{BF}_3\text{-Et}_2\text{O}$  (0.2eq) in anhydrous  $\text{CH}_2\text{Cl}_2$  was added dropwise over a period of 10 min below  $-70^\circ\text{C}$ . The reaction mixture was kept below  $-70^\circ\text{C}$  for 1hr, and then filtered. The residual molecular sieves were washed with EtOAc. The combined filtrate and washings were then worked-up by the standard method to afford a syrup. The product was purified by P-TLC developed with the following solvent combinations (v/v) to afford the corresponding glycoside: aglycon 55, EtOAc/n-hexane (1:4) three times and then EtOAc/benzene (5:95) twice; aglycon 56, EtOAc/n-hexane (1:2) and then EtOAc/benzene (5:95) twice; aglycon 57, EtOAc/n-hexane (1:4) twice and then EtOAc/benzene (5:95) twice; aglycon 66, EtOAc/n-hexane (1:2) twice and then  $\text{CH}_2\text{Cl}_2$  four times; aglycon 67, EtOAc/n-hexane (1:3) four times and then MeOH/ $\text{CH}_2\text{Cl}_2$  (1:99) twice; aglycon 68, EtOAc/n-hexane (1:4) twice; aglycon 69, EtOAc/n-hexane

(1:4) twice and then EtOAc/benzene (5:95).

**Glycoside 63** ( $\beta$ -anomer, 79.3% yield); m.p. 106-107°C;  $^1\text{H-NMR}$ : 0.89 (3H, t,  $J=8.0$ ,  $-\text{C}-\text{C}-\text{CH}_3$ ), 1.35-1.56 (2H, m,  $-\text{C}-\text{CH}_2-\text{C}$ ), 1.99 (3H, s), 2.09 (3H, s), 3.77 (3H, s,  $-\text{OCH}_3$ ), 4.06 and 4.32 (2H, two d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.41 (1H, d,  $J=8.0$ ,  $\text{C}_1-\text{H}$ ), 4.68 (1H, d,  $J=8.0$ ,  $\text{C}_1-\text{H}$ ), 5.00 (1H, dd,  $J=9.5$ , 8.0,  $\text{C}_2-\text{H}$ ), 6.78 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ), 7.06 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ );  $^{13}\text{C-NMR}$ : 96.4 ( $-\text{CCl}_3$ ), 101.4 (C-1), 102.6 (C-1'). Anal. Calcd. for  $\text{C}_{50}\text{H}_{59}\text{O}_{14}\text{Cl}_3$ : C, 60.64; H, 5.96. Found: C, 60.55; H, 5.87.

**Glycoside 64** ( $\beta$ -anomer, 72.5% yield); m.p. 100-102°C;  $^1\text{H-NMR}$ : 1.99 (3H, s), 2.09 (3H, s), 3.78 (3H, s,  $-\text{OCH}_3$ ), 4.05 and 4.33 (2H, two d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.41 (1H, d,  $J=8.0$ ,  $\text{C}_1-\text{H}$ ), 4.68 (1H, d,  $J=8.0$ ,  $\text{C}_1-\text{H}$ ), 5.19 (1H, dd,  $J=8.5$ , 8.0,  $\text{C}_2-\text{H}$ ), 5.24 (1H, dd,  $J=10.5$ , 1.5,  $-\text{C}-\text{C}=\text{CH}_{\text{cis}}$ ), 5.37 (1H, dd,  $J=17.0$ , 1.5,  $-\text{C}-\text{C}=\text{CH}_{\text{trans}}$ ), 5.82 (1H, ddt,  $J=17.0$ , 10.5, 5.3,  $-\text{C}-\text{CH}=\text{C}$ ), 6.79 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ), 7.06 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ );  $^{13}\text{C-NMR}$ : 96.5 ( $-\text{CCl}_3$ ), 101.3 (C-1), 102.7 (C-1'). Anal. Calcd. for  $\text{C}_{50}\text{H}_{57}\text{O}_{14}\text{Cl}_3$ : C, 60.75; H, 5.77. Found: C, 60.24; H, 5.75.

**Glycoside 65** ( $\beta$ -anomer, 71.7% yield);  $^1\text{H-NMR}$ : 1.98 (3H, s), 2.11 (3H, s), 2.37 (1H, t,  $J=2.5$ ,  $-\text{CH}_2-\text{C}=\text{C}$ ), 3.77 (3H, s,  $-\text{OCH}_3$ ), 4.07 and 4.33 (2H, two d,  $J=12.5$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.41 (1H, d,  $J=8.0$ ,  $\text{C}_1-\text{H}$ ), 4.72 (1H, d,  $J=8.5$ ,  $\text{C}_1-\text{H}$ ), 4.94 (1H, t,  $J=8.5$ ,  $\text{C}_2-\text{H}$ ), 6.78 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ), 7.06 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ );  $^{13}\text{C-NMR}$ : 96.4 ( $-\text{CCl}_3$ ), 101.3 (C-1), 103.1 (C-1'). Anal. Calcd. for  $\text{C}_{50}\text{H}_{55}\text{O}_{14}\text{Cl}_3$ : C, 60.88;

H, 5.58. Found: C, 59.90; H, 5.41.

**Glycoside 75** ( $\beta$ -anomer, 35.3% yield);  $^1\text{H-NMR}$ : 3.32 (1H, dd,  $J=10.0, 8.0$ ,  $\text{C}_2\text{-H}$ ), 3.82 (3H, s,  $-\text{OCH}_3$ ), 4.18 and 4.42 (2H, two d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.36 (1H, d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ), 4.94 (1H, d,  $J=7.5$ ,  $\text{C}_1\text{-H}$ ), 6.82 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ), 7.03 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ).

**Glycoside 77** ( $\beta$ -anomer, 57.0% yield);  $^1\text{H-NMR}$ : 2.09 (3H, s), 3.26 (1H, dd,  $J=9.0, 8.0$ ,  $\text{C}_2\text{-H}$ ), 3.78 (3H, s,  $-\text{OCH}_3$ ), 4.16 and 4.40 (2H, two d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.26 (1H, d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ), 4.89 (1H, d,  $J=6.5$ ,  $\text{C}_1\text{-H}$ ), 6.85 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ), 7.06 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ );  $^{13}\text{C-NMR}$ : 96.3 ( $-\text{CCl}_3$ ), 101.6 (C-1), 103.4 (C-1').

**Glycoside 78** ( $\alpha$ -anomer, 7.7% yield);  $^1\text{H-NMR}$ : 2.09 (3H, s), 3.46 (1H, dd,  $J=10.0, 3.5$ ,  $\text{C}_2\text{-H}$ ), 3.77 (3H, s,  $-\text{OCH}_3$ ), 4.16 and 4.42 (2H, two d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.88 (1H, d,  $J=7.5$ ,  $\text{C}_1\text{-H}$ ), 5.03 (1H, d,  $J=3.5$ ,  $\text{C}_1\text{-H}$ ), 6.80 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ), 7.08 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ).

**Glycoside 79** ( $\beta$ -anomer, 23.8% yield);  $^1\text{H-NMR}$ : 1.20 (9H, s), 1.21 (9H, s), 1.27 (9H, s), 3.77 (3H, s,  $-\text{OCH}_3$ ), 4.15 and 4.32 (2H, two d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.41 (1H, d,  $J=6.5$ ,  $\text{C}_1\text{-H}$ ), 4.87 (1H, d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ), 6.82 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ), 7.06 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ );  $^{13}\text{C-NMR}$ : 96.1 ( $-\text{CCl}_3$ ), 100.6 (C-1 and C-1').

**Glycoside 80** ( $\alpha$ -anomer, 55.6% yield);  $^1\text{H-NMR}$ : 1.17 (9H, s), 1.19 (9H, s), 1.20 (9H, s), 3.77 (3H, s,  $-\text{OCH}_3$ ), 4.10 and 4.27 (2H, two d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.85 (1H, d,  $J=7.0$ ,  $\text{C}_1\text{-H}$ ), 5.06 (1H, d,  $J=3.0$ ,  $\text{C}_1\text{-H}$ ), 6.78 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ), 7.03 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ );  $^{13}\text{C-NMR}$ : 95.8 (C-

1'), 96.0 ( $-\text{CCl}_3$ ), 99.8 (C-1).

Glycoside 81 ( $\beta$ -anomer, 3.2% yield);  $^1\text{H}$ -NMR: 3.78 (3H, s,  $-\text{OCH}_3$ ), 3.98 and 4.02 (2H, two d,  $J=6.5$ ,  $-\text{COCH}_2\text{Cl}$ ), 4.06 (4H, s,  $-\text{COCH}_2\text{Cl}$ ), 4.09 and 4.33 (2H, two d,  $J=11.5$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.26 (1H, d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ), 4.83 (1H, d,  $J=7.5$ ,  $\text{C}_1\text{-H}$ ), 5.13 (1H, dd,  $J=9.5$ ,  $7.5$ ,  $\text{C}_2\text{-H}$ ), 6.82 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ), 7.04 (2H, d,  $J=8.0$ ,  $\text{C}_6\text{H}_4\text{OCH}_3$ );  $^{13}\text{C}$ -NMR: 96.0 ( $-\text{CCl}_3$ ), 100.1 (C-1), 100.8 (C-1').

Glycoside 82 ( $\alpha$ -anomer, 16.2% yield);  $^1\text{H}$ -NMR: 3.46 (1H, dd,  $J=9.5$ ,  $3.5$ ,  $\text{C}_2\text{-H}$ ), 3.74 and 3.80 (2H, two d,  $J=7.0$ ,  $-\text{COCH}_2\text{Cl}$ ), 4.07 (2H, s,  $-\text{COCH}_2\text{Cl}$ ), 4.08 (2H, s,  $-\text{COCH}_2\text{Cl}$ ), 4.13 and 4.39 (2H, two d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.80 (1H, d,  $J=3.5$ ,  $\text{C}_1\text{-H}$ ), 4.85 (1H, d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ), 5.35 (1H, dd,  $J=9.0$ ,  $8.0$ ,  $\text{C}_2\text{-H}$ ), 6.82 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ), 7.04 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ).



#### 4. EXPERIMENTAL (Chapter 3.)

##### 2,4,6-tri-O-acetyl-3-O-benzyl-D-glucopyranose (91)

A solution of compound 25 (300 mg, 0.57 mM) in  $\text{Ac}_2\text{O}$  (5 ml) was treated with zinc powder (930 mg, 14.2 mM) at room temperature for 2 h. The reaction mixture was filtered and washed with EtOAc. The filtrate and washings were worked-up by the standard method to give a colorless oil. The oil was purified by P-TLC (EtOAc / *n*-hexane (1:2)) to give compound 91 as a oil (154 mg, 68.3% yield).

##### 2,2,2-Trichloroethyl 3-O-benzyl-4,6-O-p-methoxybenzylidene- $\beta$ -D-glucopyranoside (92)

To a solution of compound 45 (10g, 32mM) and p-methoxybenzaldehyde dimethylacetal (11.7 ml, 64 mM) in dioxane (100 ml), p-toluenesulfonic acid (400 mg) was added. The reaction mixture was kept at 50°C under 15 mmHg for 20 min, and then worked-up by the standard method to give a colorless oil. The oil was crystallized from EtOH / *iso*-PrOH / *n*-hexane to afford colorless crystals (11.8 g, 85.5% yield).

##### 2,2,2-Trichloroethyl 2-O-acetyl-3-O-benzyl-4-O-p-methoxybenzyl- $\beta$ -D-glucopyranoside (95) and 2,2,2-trichloroethyl 2-O-acetyl-3-O-benzyl-6-O-p-methoxybenzyl- $\beta$ -D-glucopyranoside (97)

Compound 93 (362mg, 0.64mM) and powdered molecular sieves 4A (200mg) were dried over  $\text{P}_2\text{O}_5$  in a vacuum

desiccator before use. To a stirred suspension of compound 93 and molecular sieves 4A in anhydrous  $\text{CH}_3\text{CN}$  (5ml),  $\text{NaCNBH}_4$  (243mg, 3.87mM) was added and then TMS-Cl (491  $\mu\text{l}$ , 3.87mM) in anhydrous  $\text{CH}_3\text{CN}$  (1ml) was added dropwise. The reaction mixture was kept at room temperature for 30 min, filtered and washed with EtOAc. The combined filtrate and washings were worked-up by the standard method to give a colorless oil. The oil was purified by P-TLC ( $\text{MeOH}/\text{CH}_2\text{Cl}_2$  (1:99)) to afford compounds 95 (140mg, 38.5% yield) and 97 (109mg, 30.0% yield) as syrups.

2,2,2-Trichloroethyl 3-O-benzyl-6-O-p-methoxybenzyl-2-O-pivaloyl- $\beta$ -D-glucopyranoside (98)

Compound 94 (121mg, 0.2mM) and powdered molecular sieves 4A (100mg) were dried over  $\text{P}_2\text{O}_5$  in a vacuum desiccator before use. To a stirred mixture containing compound 94 and molecular sieves 4A in anhydrous  $\text{CH}_3\text{CN}$  (2.5ml),  $\text{NaCNBH}_4$  (75.6mg, 1.2mM) was added and then TMS-Cl (153  $\mu\text{l}$ , 1.2mM) in anhydrous  $\text{CH}_3\text{CN}$  (0.5mM) was added dropwise. After 3 h, the reaction did not complete yet, and then TMS-Cl (76  $\mu\text{l}$ , 0.6mM) in anhydrous  $\text{CH}_3\text{CN}$  (0.5ml) was added dropwise. The reaction mixture was kept for another 30 min at room temperature, filtered and worked-up by the standard method to yield a colorless oil. The oil was purified by P-TLC ( $\text{MeOH}/\text{CH}_2\text{Cl}_2$  (5:95)) to afford compound 98 (73.2mg, 60.3% yield) as a syrup;  $^1\text{H-NMR}$ : 1.23 (9H, s), 3.46-3.81 (5H, m,  $\text{C}_3\text{-H}$ ,  $\text{C}_4\text{-H}$ ,  $\text{C}_5\text{-H}$ ,  $\text{C}_6\text{-H}_a$ ,  $\text{C}_6\text{-H}_b$ ), 3.83 (3H, s,  $-\text{OCH}_3$ ), 4.19 and 4.36 (2H, two d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.51 and 4.59 (2H, two d,  $J=12.0$ ,

benzyl), 4.72 and 4.79 (2H, two d,  $J=12.0$ , benzyl), 4.80 (1H, d,  $J=8.0$ ,  $C_1$ -H), 5.14 (1H, dd,  $J=10.0$ ,  $8.0$ ,  $C_2$ -H), 6.92 (2H, d,  $J=8.0$ ,  $-C_6H_4OCH_3$ ), 7.29 (2H, d,  $J=8.0$ ,  $-C_6H_4OCH_3$ ), 7.30-7.44 (5H, m, phenyl).

**2,3,6-tri-O-benzyl-4-O-monochloroacetyl-D-glucopyranose**  
(112)

A mixture of compound 111 (519mg, 0.92mM),  $SeO_2$  (111mg, 1.0mM) and AcOH (57  $\mu$ l, 1.0mM) in dioxane (5ml) was heated under reflux for 2hr. The reaction mixture was worked-up by standard method to give a brown oil. The oil was purified by P-TLC (EtOAc/ $n$ -hexane (1:3)) to afford compound 112 (298mg, 61.9% yield) as crystals; m.p. 104-105°C;  $^1H$ -NMR : 3.37-3.63 (2H, m,  $C_6-H_a$ ,  $C_6-H_b$ ), 3.52 and 3.63 (2H, two d,  $J=14.5$ ,  $-COCH_2Cl$ ), 3.62 (1H, dd,  $J=9.5$ ,  $3.5$ ,  $C_2$ -H), 3.93 (1H, t,  $J=9.5$ ,  $C_3$ -H), 4.09 (1H, dt,  $J=9.5$ ,  $3.5$ ,  $C_5$ -H), 4.42 and 4.51 (2H, two d,  $J=11.5$ , benzyl), 4.63 and 4.88 (2H, two d,  $J=11.5$ , benzyl), 4.68 and 4.79 (2H, two d,  $J=11.5$ , benzyl), 5.06 (1H, dd,  $J=9.5$ ,  $9.5$ ,  $C_4$ -H), 5.19 (1H, d,  $J=3.5$ ,  $C_1$ -H), 7.20-7.44 (15H, m, phenyl).

**Imidoylation of compounds 91,102,103,105,109,112,116 and 117**

To a solution of the respective compound (1.0eq) in anhydrous  $CH_2Cl_2$ , DBU (0.2eq) and  $CCl_3CN$  (2.0eq) were added. The reaction solution was stirred for 3hr at room temperature, and then evaporated below 25°C to give a yellow oil. The oil was purified by P-TLC developed with the

following solvent combinations (v/v) to give the corresponding imidate: compounds 91,102,105,109 and 116, EtOAc/n-hexane (1:2) ; compounds 103,112 and 117, EtOAc/n-hexane (1:4).

Imidate 83;  $^1\text{H-NMR}$ : 1.96 (3H, s), 2.05 (3H, s), 3.68 (1H, t,  $J=9.5$ ,  $\text{C}_4\text{-H}$ ), 3.82 (3H, s,  $-\text{OCH}_3$ ), 3.97-4.11 (1H, m,  $\text{C}_5\text{-H}$ ), 4.11 (1H, t,  $J=9.5$ ,  $\text{C}_3\text{-H}$ ), 4.18-4.40 (2H, m,  $\text{C}_6\text{-H}_a$ ,  $\text{C}_6\text{-H}_b$ ), 4.53 and 4.80 (2H, two d,  $J=10.5$ , benzyl), 4.79 and 4.89 (2H, two d,  $J=11.0$ , benzyl), 5.04 (1H, dd,  $J=9.5$ , 4.0,  $\text{C}_2\text{-H}$ ), 6.44 (1H, d,  $J=4.0$ ,  $\text{C}_1\text{-H}$ ), 6.85 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ), 7.20 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ), 7.25-7.41 (5H, m, phenyl), 8.58 (1H, s,  $-\text{C}(=\text{NH})\text{CCl}_3$ ).

Imidate 84;  $^1\text{H-NMR}$ : 1.18 (9H, s), 1.21 (9H, s), 3.67 (1H, t,  $J=9.5$ ,  $\text{C}_4\text{-H}$ ), 3.82 (3H, s,  $-\text{OCH}_3$ ), 4.02-4.16 (1H, m,  $\text{C}_5\text{-H}$ ), 4.14 (1H, t,  $J=9.5$ ,  $\text{C}_3\text{-H}$ ), 4.24 (1H, dd,  $J=12.0$ , 4.0,  $\text{C}_6\text{-H}_a$ ), 4.38 (1H, dd,  $J=12.0$ , 2.0,  $\text{C}_6\text{-H}_b$ ), 4.49 and 4.78 (2H, two d,  $J=10.0$ , benzyl), 4.82 and 4.87 (2H, two d,  $J=11.5$ , benzyl), 5.06 (1H, dd,  $J=10.0$ , 3.5,  $\text{C}_2\text{-H}$ ), 6.52 (1H, d,  $J=3.5$ ,  $\text{C}_1\text{-H}$ ), 6.87 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ), 7.20 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ), 7.26-7.48 (5H, m, phenyl), 8.62 (1H, s,  $-\text{C}(=\text{NH})\text{CCl}_3$ ).

Imidate 85;  $^1\text{H-NMR}$  : 3.44 (1H, dd,  $J=10.0$ , 3.5,  $\text{C}_6\text{-H}_a$ ), 3.53 (1H, dd,  $J=10.0$ , 3.5,  $\text{C}_6\text{-H}_b$ ), 3.56 and 3.64 (2H, two d,  $J=14.5$ ,  $-\text{COCH}_2\text{Cl}$ ), 3.81 (1H, dd,  $J=9.5$ , 3.5,  $\text{C}_2\text{-H}$ ), 3.97 (1H, t,  $J=9.5$ ,  $\text{C}_3\text{-H}$ ), 4.01 (1H, dt,  $J=10.0$ , 3.5,  $\text{C}_5\text{-H}$ ), 4.40 and 4.50 (2H, two d,  $J=12.0$ , benzyl), 4.62 and 4.87 (2H, two d,  $J=12.0$ , benzyl), 4.68 and 4.75 (2H, two d,  $J=11.5$ , benzyl), 5.23 (1H, dd,  $J=10.0$ , 9.5,  $\text{C}_4\text{-H}$ ), 6.49 (1H, d,

$J=3.5$ ,  $C_1-H$ ), 7.18-7.42 (15H, m, phenyl), 8.64 (1H, s,  $-C(=NH)CCl_3$ ).

Imidate 86;  $^1H-NMR$  : 1.96 (3H, s), 2.13 (3H, s), 6.61 (1H, d,  $J=3.5$ ,  $C_1-H$ ), 8.74 (1H, s,  $-C(=NH)CCl_3$ ).

Imidate 87;  $^1H-NMR$  : 1.20 (9H, s), 1.22 (9H, s), 3.66 and 3.85 (2H, two d,  $J=15.0$ ,  $-COCH_2Cl$ ), 4.02-4.24 (4H, m,  $C_3-H$ ,  $C_5-H$ ,  $C_6-H_a$ ,  $C_6-H_b$ ), 4.57 and 4.81 (2H, two d,  $J=11.5$ , benzyl), 5.08 (1H, dd,  $J=10.0$ , 3.5,  $C_2-H$ ), 5.19 (1H, t,  $J=10.0$ ,  $C_4-H$ ), 6.60 (1H, d,  $J=3.5$ ,  $C_1-H$ ), 7.22-7.46 (5H, m, phenyl), 8.72 (1H, s,  $-C(=NH)CCl_3$ ).

Imidate 88;  $^1H-NMR$  : 1.91 (3H, s), 1.99 (3H, s), 3.42-3.58 (2H, m,  $C_6-H_a$ ,  $C_6-H_b$ ), 3.78 (3H, s,  $-OCH_3$ ), 3.98-4.16 (1H, m,  $C_5-H$ ), 4.05 (1H, t,  $J=9.5$ ,  $C_3-H$ ), 4.39 and 4.46 (2H, two d,  $J=12.0$ , benzyl), 4.62 and 4.71 (2H, two d,  $J=12.0$ , benzyl), 5.07 (1H, dd,  $J=9.5$ , 3.5,  $C_2-H$ ), 5.23 (1H, t,  $J=9.5$ ,  $C_4-H$ ), 6.56 (1H, d,  $J=3.5$ ,  $C_1-H$ ), 6.88 (2H, d,  $J=8.0$ ,  $-C_6H_4OCH_3$ ), 7.20-7.40 (6H, m, phenyl), 8.63 (1H, s,  $-C(=NH)CCl_3$ ).

Imidate 89;  $^1H-NMR$  : 1.96 (3H, s), 2.05 (3H, s), 3.79 (1H, dd,  $J=10.0$ , 4.0,  $C_2-H$ ), 3.97 (1H, t,  $J=10.0$ ,  $C_3-H$ ), 4.00-4.13 (1H, m,  $C_5-H$ ), 4.06 (1H, dd,  $J=13.0$ , 2.5,  $C_6-H_a$ ), 4.21 (1H, dd,  $J=13.0$ , 5.0,  $C_6-H_b$ ), 4.66 and 4.88 (2H, two d,  $J=12.0$ , benzyl), 4.69 and 4.75 (2H, two d,  $J=12.5$ , benzyl), 5.10 (1H, t,  $J=10.0$ ,  $C_4-H$ ), 6.45 (1H, d,  $J=4.0$ ,  $C_1-H$ ), 7.24-7.33 (10H, m, phenyl), 8.64 (1H, s,  $-C(=NH)CCl_3$ ).

Imidate 90;  $^1H-NMR$  : 2.00 (3H, s), 2.01 (3H, s), 2.09 (3H, s), 3.95-4.16 (1H, m,  $C_5-H$ ), 4.09 (1H, dd,  $J=13.0$ , 2.5,

$C_6-H_a$ ), 4.12 (1H, t,  $J=10.0$ ,  $C_3-H$ ), 4.21 (1H, dd,  $J=13.0$ , 5.0,  $C_6-H_b$ ), 4.64 and 4.73 (2H, two d,  $J=11.5$ , benzyl), 5.08 (1H, dd,  $J=10.0$ , 4.0,  $C_2-H$ ), 5.20 (1H, t,  $J=10.0$ ,  $C_4-H$ ), 6.54 (1H, d,  $J=4.0$ ,  $C_1-H$ ), 7.22-7.40 (5H, m, phenyl), 8.67 (1H, s,  $-C(=NH)CCl_3$ ).

#### Glycosylation between imidates 34,83-90 and aglycon 40

The respective imidate (1.0eq) and aglycon 40 (2.0eq) were dried over  $P_2O_5$  in a vacuum desiccator; powdered molecular sieves 4A were dried overnight in a 120 °C oven before use. A solution of imidate and aglycon in anhydrous  $CH_2Cl_2$  was added to a stirred suspension of molecular sieves 4A in anhydrous  $CH_2Cl_2$  below -70°C. To this reaction mixture,  $BF_3-Et_2O$  (0.2eq) in anhydrous  $CH_2Cl_2$  was added dropwise below -70°C. The reaction mixture was then kept below -70°C for 1 hr, filtered and washed with EtOAc. The combined filtrate and washings were worked-up by the standard method to give a colorless oil. The oil was purified by P-TLC developed with the following solvent combinations (v/v) to afford the corresponding glycoside : imidate 34, EtOAc / n-hexane (1:4); imidate 83, MeOH /  $CH_2Cl_2$  (2:98); imidate 84, EtOAc / n-hexane (1:2), imidate 85,  $CH_2Cl_2$ .

Glycoside 53 ( $\beta$ -anomer, 82.0% yield).

Glycoside 118 ( $\beta$ -anomer, 60.0% yield);  $^1H$ -NMR : 1.90 (3H, s), 1.91 (3H, s), 1.98 (3H, s), 2.13 (3H, s), 3.81 (3H, s,  $-OCH_3$ ), 4.07 and 4.32 (2H, two d,  $J=12.0$ ,  $-OCH_2CCl_3$ ), 4.52 (1H, d,  $J=8.0$ ,  $C_1-H$ ), 4.68 (1H, d,  $J=8.0$ ,  $C_1-H$ ), 6.87

(2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ), 7.19 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ), 7.20-7.46 (10H, m, phenyl).

**Glycoside 119** ( $\beta$ -anomer, 57.8% yield);  $^1\text{H-NMR}$  : 1.16 (9H, s), 1.20 (9H, s), 1.89 (3H, s), 2.14 (3H, s), 3.81 (3H, s,  $-\text{OCH}_3$ ), 4.07 and 4.34 (2H, two d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.46 (1H, d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ), 6.86 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ), 7.16 (2H, d,  $J=8.0$ ,  $-\text{C}_6\text{H}_4\text{OCH}_3$ ), 7.22-7.44 (10H, m, phenyl).

**Glycoside 120** ( $\beta$ -anomer, 13.5% yield);  $^1\text{H-NMR}$  : 1.98 (3H, s), 2.05 (3H, s), 3.48 and 3.59 (2H, two d,  $J=14.5$ ,  $-\text{COCH}_2\text{Cl}$ ), 4.07 and 4.33 (2H, d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.43 (1H, d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ), 4.69 (1H, d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ), 7.18-7.40 (20H, m, phenyl).

**Glycoside 121** ( $\alpha$ -anomer, 3.0% yield);  $^1\text{H-NMR}$  : 1.96 (3H, s), 2.05 (3H, s), 3.54 and 3.64 (2H, two d,  $J=14.5$ ,  $-\text{COCH}_2\text{Cl}$ ), 3.55 (1H, dd,  $J=9.5$ , 3.5,  $\text{C}_2\text{-H}$ ), 4.09 and 4.37 (2H, two d,  $J=12.0$ ,  $-\text{OCH}_2\text{CCl}_3$ ), 4.76 (1H, d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ), 5.39 (1H, d,  $J=3.5$ ,  $\text{C}_1\text{-H}$ ), 7.10-7.40 (20H, m, phenyl).

**Allyl 2,4,6-tri-O-acetyl-3-O-benzyl- $\beta$ -D-glucopyranoside (124)**

Compound **44** (1000mg, 2.28mM) was dried over  $\text{P}_2\text{O}_5$  in a vacuum desiccator. To a solution of compound **44** in anhydrous  $\text{CH}_2\text{Cl}_2$  (10 ml), allyl alcohol (0.78ml, 11.4mM) was added, and then  $\text{BF}_3\text{-Et}_2\text{O}$  (2.8ml, 22.8mM) was added dropwise at  $0^\circ\text{C}$ . The reaction solution was kept at  $0^\circ\text{C}$  for 4 h. The reaction mixture was worked-up by the standard method to give a yellow oil. The product was crystallized from EtOH /  $n$ -

hexane to afford **124** (770mg, 77.4%); m.p. 76-77°C;  $^1\text{H-NMR}$ : 1.99 (s, 3H), 2.03 (s, 3H), 2.10 (s, 3H), 3.58 (1H, ddd,  $J=9.5$ , 5.0, 2.5,  $\text{C}_5\text{-H}$ ), 3.70 (1H, t,  $J=9.5$ ,  $\text{C}_3\text{-H}$ ), 4.07 (1H, ddd,  $J=13.5$ , 6.0, 1.5,  $-\text{CH}_2\text{-C}=\text{C}$ ), 4.11 (1H, dd,  $J=12.0$ , 2.5,  $\text{C}_6\text{-H}_a$ ), 4.22 (1H, dd,  $J=12.0$ , 5.0,  $\text{C}_6\text{-H}_b$ ), 4.33 (1H, ddd,  $J=13.5$ , 5.0, 1.5,  $-\text{CH}_2\text{-C}=\text{C}$ ), 4.46 (1H, d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ), 4.57 (1H, d,  $J=12.0$ , benzyl), 4.63 (1H, d,  $J=12.0$ , benzyl), 5.09 (1H, dd,  $J=9.5$ , 8.0,  $\text{C}_2\text{-H}$ ), 5.13 (1H, t,  $J=9.5$ ,  $\text{C}_4\text{-H}$ ), 5.18 (1H, dd,  $J=10.0$ , 1.5,  $-\text{C}=\text{C}=\text{CH}_{\text{cis}}$ ), 5.26 (1H, dd,  $J=17.5$ , 1.5,  $-\text{C}=\text{C}=\text{CH}_{\text{trans}}$ ), 5.84 (1H, dddd,  $J=17.5$ , 10.0, 6.0, 5.0,  $-\text{C}=\text{CH}=\text{C}$ ), 7.20-7.40 (5H, m, phenyl). Anal. Calc. for  $\text{C}_{22}\text{H}_{28}\text{O}_9$ : C, 60.55; H, 6.42. Found: C, 59.75; H, 6.61.

**Allyl 3-O-benzyl- $\beta$ -D-glucopyranoside (125)**

Compound **125** was prepared from compound **124** in the same way that compound **45** was prepared from compound **25**, (28% NaOMe in MeOH / MeOH, at 50°C for 1 h, 93.8% yield).

**Allyl 3-O-benzyl-4,6-O-p-methoxybenzylidene- $\beta$ -D-glucopyranoside (126)**

Compound **126** was obtained from compound **125** in the same way that compound **92** was obtained from compound **45**, ( $p\text{-MeOPhCH(OMe)}_2$  /  $p\text{-TsOH}$  / dioxane, at 50°C under 15 mmHg, for 20 min, 85.8% yield).

**Allyl 3-O-benzyl-4-O-p-methoxybenzyl- $\beta$ -D-glucopyranoside (127)**

To a solution of compound **126** (3.26 g, 7.6 mM) in



anhydrous  $\text{CH}_2\text{Cl}_2$  (40 ml) and anhydrous  $\text{Et}_2\text{O}$  (20 ml),  $\text{LiAlH}_4$  (1.45 g, 38.1 mM) was added. A solution of  $\text{AlCl}_3$  (4.06 g, 30.4 mM) in anhydrous  $\text{Et}_2\text{O}$  (20 ml) was added dropwise at  $0^\circ\text{C}$  over 30 min. The reaction mixture was stirred at  $0^\circ\text{C}$  for 1 h, treated with  $\text{H}_2\text{O}$  (2.6 ml) in tetrahydrofuran (10 ml) to decompose the excess reagents and then filtered with  $\text{EtOAc}$ . The combined filtrate and washings were worked-up by the standard method to give a colorless syrup. The product was crystallized from  $\text{EtOH}$  / *n*-hexane to afford colorless crystals (2.74 g, 83.6% yield).

Allyl 2,6-di-O-acetyl-3-O-benzyl-4-O-p-methoxybenzyl- $\beta$ -D-glucopyranoside (122)

Compound 122 was synthesized from compound 127 by the standard acetylation method, ( $\text{Ac}_2\text{O}$  / pyridine, at  $50^\circ\text{C}$ , for 2 h, a quantitative yield);  $^1\text{H-NMR}$ : 1.99 (3H, s), 2.06 (3H, s), 3.40-3.55 (1H, m,  $\text{C}_5\text{-H}$ ), 3.56-3.73 (2H, m,  $\text{C}_3\text{-H}$ ,  $\text{C}_5\text{-H}$ ), 3.79 (3H, s,  $-\text{OCH}_3$ ), 4.04 (1H, dd,  $J=13.5$ , 6.0,  $-\text{CH}_2\text{-C}=\text{C}$ ), 4.20 (1H, dd,  $J=12.0$ , 4.0,  $\text{C}_6\text{-H}_a$ ), 4.31 (1H, dd,  $J=13.5$ , 4.0,  $-\text{CH}_2\text{-C}=\text{C}$ ), 4.34 (1H, dd,  $J=12.0$ , 2.0,  $\text{C}_6\text{-H}_b$ ), 4.40 (1H, d,  $J=8.0$ ,  $\text{C}_1\text{-H}$ ), 4.49 (1H, d,  $J=10.5$ , benzyl), 4.69 (1H, d,  $J=11.0$ , benzyl), 4.76 (1H, d,  $J=10.5$ , benzyl), 4.82 (1H, d,  $J=11.0$ , benzyl), 5.01 (1H, dd,  $J=9.0$ , 8.0,  $\text{C}_2\text{-H}$ ), 5.17 (1H, dd,  $J=10.5$ , 1.5,  $-\text{C}=\text{C}=\text{CHcis}$ ), 5.24 (1H, dd,  $J=17.0$ , 1.5,  $-\text{C}=\text{C}=\text{CHtrans}$ ), 5.74-5.96 (1H, m,  $-\text{C}-\text{CH}=\text{C}$ ), 6.88 (2H, d,  $J=8.0$ ,  $-\text{OC}_6\text{H}_4\text{OCH}_3$ ), 7.22 (2H, d,  $J=8.0$ ,  $-\text{OC}_6\text{H}_4\text{OCH}_3$ ), 7.28-7.44 (5H, m, phenyl).

Allyl 2,6-di-O-pivaloyl-3-O-benzyl-4-O-p-methoxybenzyl- $\beta$ -D-glucopyranoside (123)

Compound 123 was obtained from compound 127 in the same way that compound 71 was obtained from compound 42, ((CH<sub>3</sub>)<sub>3</sub>CCOCl / pyridine / 50°C / 12 h, a quantitative yield).

2,6-Di-O-acetyl-3-O-benzyl-4-O-p-methoxybenzyl-D-glucopyranoside (102)

A mixture of compound 122 (252 mg, 0.49 mM), SeO<sub>2</sub> (60 mg, 0.54 mM) and AcOH (31  $\mu$ l, 0.54 mM) in dioxane (2 ml) was heated under reflux for 3 h. The reaction mixture was worked-up by the standard method to give a yellow oil. The oil was purified by P-TLC (EtOAc / n-hexane (1:1)) to afford compound 102 (140 mg, 60.3% yield).

Allyl 2,6-di-O-acetyl-3-O-benzyl- $\beta$ -D-glucopyranoside (128)

To a solution of compound 122 (372 mg, 0.72 mM) in CH<sub>3</sub>CN-H<sub>2</sub>O (4 ml, 9:1), Ce(NH<sub>4</sub>)<sub>2</sub>(NO<sub>3</sub>)<sub>6</sub> (1.19 g, 2.16 mM) was added. The reaction mixture was kept at room temperature for 15 min, and then worked-up by the standard method to give a colorless oil. The oil was purified by P-TLC (EtOAc / n-hexane (1:2)) to afford a colorless oil (243 mg, 85.3% yield).

## REFERENCES

- 1) Abbas, S.; Haines, A.H.: Use of the 2,6-dimethoxybenzoyl group to prevent acyl migration during Purdie methylation, a synthesis of 2-O-methyl-D-mannose, Carbohydr. Res., 41, 298-303 (1975).
- 2) Bochkov, A.F. and Zaikov, G.E.: "Chemistry of the O-glycosidic bond.", translated by Schuerch, C., Pergamon Press, New York, (1979).
- 3) Boeckel, C.A.A.; Beetz, T. and Aelst, S.F.: Substituent Effects on Carbohydrate Coupling Reactions Promoted by Insoluble Silver Salts, Tetrahedron, 40, 4097-4107 (1984).
- 4) Boeckel, C.A.A. and Beetz, T.: Substituent Effects in Carbohydrate Chemistry Part II, Coupling reactions involving gluco- and galacto-pyranosyl bromides promoted by insoluble silver salts, Recl. Trav. Chim. Pays-Bas, 104, 171-173 (1985).
- 5) Boeckel, C.A.A. and Beetz, T.: Substituent Effects in Carbohydrate Chemistry Part III, Effects of C-2 substituents on coupling reactions promoted by insoluble silver salts, Recl. Trav. Chim. Pays-Bas, 104, 174-176 (1985).
- 6) Bonner, W.A.: On the structure of 1,2,3,6-tetra-O-acetyl- $\beta$ -D-glucose, J. Am. Chem. Soc., 80, 3697-3700, (1958).
- 7) Charton, M.: "Progress in Physical Organic Chemistry", John Willey & Sons, New York, 1981, p 119.
- 8) Chaudhary, S.K. and Hernandez, O.: A Simplified Procedure

- for The Preparation of Triphenylmethyl Ethers,  
Tetrahedron Letters, No.2, 95-98 (1979).
- 9) Cook, A.F. and Maichnk, D.T.: Use of chloroacetic anhydride for the protection of nucleoside hydroxyl groups, J. Org. Chem., 55, 1940-1943 (1970).
- 10) Corey, E.J.; Gras, J.L. and Ulrich, P.: A New General Method for Protection of The Hydroxyl Function, Tetrahedron Letters, No.11, 809-812 (1976).
- 11) Cram, J.D. and Allinger, L.N.: Macro Rings XIII, Synthesis and properties of 1,7-cyclododecadiyne and related compounds, J. Am. Chem. Soc., 78, 2518-2524 (1956).
- 12) Cramer, F.; Pawelzik, K. and Baldauf, H.J.: Darstellung von Trichloroacetimidsäurerstern, Chem. Ber., 91, 1049-1054 (1958).
- 13) Dahmen, J.; Frejd, T.; Magnusson, G. and Noori, G.: Boron trifluoride etherate induced glycosylation, formation of alkyl glycosides and thioglycosides of 2-deoxy-2-phthalimidoglycopyranoses, Carbohydr. Res., 114, 328-330 (1983).
- 14) Dewar, M.J.S.: "The Electron Theory of Organic Chemistry", Clarendon Press, Oxford, 1949, p 168.
- 15) Dmitriev, B.A.; Nikolaev, A.V.; Shashkov, A.S. and Kochetkov, N.K.: Block-synthesis of higher oligosaccharides, Synthesis of hexa- and nona-saccharide fragments of the O-antigenic polysaccharide of Salmonella newington, Carbohydr. Res., 100, 195-206

(1982).

- 16) Evans, M.E.: Methyl 4,6-O-benzylidene- $\alpha$ -and- $\beta$ -D-glucosides, Carbohydr. Res., 21, 473-475 (1972).
- 17) Franchimont, A.P.N.: Uber Kohlehydrate, Ber., 12, 1938-1942 (1879).
- 18) Frechet, J.M. and Schuerch, C.: Solid-phase synthesis of oligosaccharides I, Preparation of the solid support, Poly[(p-propen-3-ol-yl) styrene], J. Am. Chem. Soc., 93, 492-496 (1971).
- 19) Freudenberg, K. and Nagai, W.: Die synthese der cellobiose, Ber., 66, 27-29 (1933).
- 20) Fujii, K.; Nakano, S. and Fujita, E.: An improved method for methoxymethylation of alcohols under mild acidic conditions, Synthesis, 276-277 (1975).
- 21) Fujimaki, I. and Kuzuhara, H.: Conversion of 1,6-Anhydromaltose into Psudodisaccharides Containing Aminocyclitols as Constituent, Agric. Biol. Chem., 44, 2055-2059 (1980).
- 22) Fukami, H.; Ikeda, S.; Kitahara, K and Nakajima, M.: Total Synthesis of Ribostamycin, Agric. Biol. Chem., 41, 1689-1694 (1977).
- 23) Gagnaire, D.Y. and Vottero, P.J.A.: Synthèse d'oligosaccharides sur polymere support Partie III, Possibilites d' utilistation du groupe chloroacetyle pour la protection de la position 6 de derives du D-glucose, Carbohydr. Res., 28, 165-170 (1973).
- 24) Garegg, P.J. and Hultberg, H.: A novel reductive ring-opening of carbohydrate benzylidene acetals with

- unusual regioselectivity, Carbohydr. Res., 93, C10-11 (1981).
- 25) Garegg, P.J.; Hultberg, H. and Wallin, S.: A novel reductive ring-opening of carbohydrate benzylidene acetals, Carbohydr. Res., 101, 97-101 (1982).
- 26) Gigg, J. and Gigg, R.: The allyl ether as protecting group in carbohydrate chemistry, J. Chem. Soc., 82-86 (1966).
- 27) Goldstein, I.J. and Hullar, T.L.: "Adv. Carbohydr. Chem. Biochem. Vol.21", Academic Press, New York and London, 1966, p 431.
- 28) Greene, T.W.: "Protective Groups in Organic Synthesis", John Wiley & Sons Inc, New York, 1981.
- 29) Haines, A.H.: "Adv. Carbohydr. Chem. Biochem. Vol.33", Academic Press, New York and London, 1976, p 100.
- 30) Haines, A.H.: "Adv. Carbohydr. Chem. Biochem. Vol.39", Academic Press, New York and London, 1981, p 13.
- 31) Hall, D.M. and Lawler, T.E.: New routes to the synthesis of 2,3,6-tri-Q-substituted methyl  $\beta$ -D-glucopyranoside, An improved synthesis of  $\alpha$ -cellotriase hendecaacetate, Carbohydr. Res., 16, 1-7 (1971).
- 32) Hall, D.M.; Lawler, T.E. and Childress, B.C.: A practical synthesis of 1,2,3,6-tetra-Q-acetyl- $\alpha$ -and  $\beta$ -D-glucopyranose and their use to prepare trisaccharides, Carbohydr. Res., 38, 359-363 (1974).
- 33) Helferich, B.; Bredereck, H. and Schneidmüller, A.: Acylwanderung an partiell acylierten methyl-glucosiden,

- Ann, 458, 111-116 (1927).
- 34) Helferich, B.: "Adv. Carbohydr. Chem. Biochem. Vol.3", Academic Press, New York, 1948, p 79.
- 35) Higuchi, T.: "Biosynthesis and Biodegradation of Wood Components", Academic Press Inc, Orlando, Florida, 1985.
- 36) Hirano, I.; Itoh, K.; Nishino, S.; Araki, Y and Ishido, Y: Regioselective phenylcarbamoylation of purine and pyrimidine ribonucleosides through dibutyltin oxide- and tertiary amine-phenyl isocyanate systems, Carbohydr. Res., 152, 159-172 (1986).
- 37) Hirano, S.: The preparation of cellulose like polymer from 2,3,6-tri-O-(N-phenylcarbamoyl)-D-glucopyranose by the action of phosphorous pentaoxide in dimethyl sulfoxide, Agric. Biol. Chem., 37, 187-189 (1973).
- 38) Hori, R. and Koizumi, T.: Studies on Alkyl Derivatives of Carbohydrate III, Syntheses of Glucose 3-Fatty Acid Monoesters, Yakugaku Zasshi, 78, 1003-1005 (1958).
- 39) Husemann, E. and Muller, G.J.: Uber die synthese unverzweigter polysaccharide, Makromol. Chem., 91, 212-230 (1966).
- 40) Igarashi, K: "Adv. Carbohydr. Chem. Biochem. Vol.34", Academic Press, New York and London, 1977, p 243.
- 41) Inversen, T. and Bundle, D.R.: Benzyl Trichloroacetimidate, A Versatile Reagent for Acid-catalysed Benzylation of Hydroxy-groups, J. Chem. Soc. Chem. Comm., 1240-1241 (1981).
- 42) Johansson, R. and Samuelsson, B.: Regioselective reductive ring-opening of 4-methoxybenzylidene acetal of

- hexopyranosides, Access to a novel protecting-group strategy Part I, J. Chem. Soc., 2371-2374 (1984).
- 43) Kawada, T.; Nakatsubo, F. and Murakami, K.: Synthetic Studies of Cellulose III, Synthesis of allyl 2,3,6-tri-O-benzyl-4-O-(p-methoxybenzyl)- $\beta$ -D-glucopyranose and its conversion into glycosyl acceptor and glycosyl donor, Mokuzai Gakkaishi, 35, 14-20 (1989).
- 44) Kawada, T.; Nakatsubo, F. and Murakami, K.: Synthetic Studies of Cellulose IV, Preparation of perbenzylated allyl 4'-O-p-methoxybenzyl cellobioside by the imidate method, Mokuzai Gakkaishi, 35, 21-25 (1989).
- 45) Kariyone, K. and Yazawa, H.: Oxidative cleavage of  $\alpha$ ,  $\beta$ -unsaturated ether, Tetrahedron Letters, No.33, 2885-2888 (1970).
- 46) Khorlin, A.Y.; Nesmeyanov, V.A. and Zurabyan, S.E.: Glycosylation of sugar 2,3-diphenyl-2-cyclopropen-1-yl ethers, A new route to oligosaccharides, Carbohydr. Res., 43, 69-77 (1975).
- 47) Kohli, V.; Blocker, H. and Koster, H.: The Triphenylmethyl (Trityl) Group And Its Uses In Nucleoside Chemistry, Tetrahedron Letters, 21, 2683-2686 (1980).
- 48) Koto, S; Morishima, S. and Zen, S.; Recent Advances in O-Glycosylation, Yuki Gosei Kagaku, 41, 701-717 (1983).
- 49) Kovac, P. and Hirsch, J.: Systematic sequential synthesis of (1-4)- $\beta$ -D-xylo-oligosaccharides and their methyl  $\beta$ -glycoside, Carbohydr. Res., 90, C5-7 (1981).



- 50) Kunz, H. and Harreus, A.: Glycosidsynthese mit 2,3,4,6-tetra-Q-pivaloyl- $\alpha$ -D-glucopyranosylbromid, Liebigs. Ann. Chem., 41-48 (1982).
- 51) Kuzuhara, H. and Ichikawa, Y.: Bioorganic Chemistry on Heparin, Nippon Nogei Kagaku Kaishi, 63, 1238-1242 (1989).
- 52) Lapatsanis, L.: A New Method for the Preparation of Diphenylmethyl Ethers, Tetrahedron Letters, No.41, 3943-3944 (1978).
- 53) Lee, R.T. and Lee, Y.C.: Synthesis of 3-(2-Aminoethylthio)propyl Glycosides, Carbohydr. Res., 37, 193-201 (1974).
- 54) Lehrfeld, J.: Silica Gel Catalyzed Detritylation of Some Carbohydrate Derivatives, J. Org. Chem., 32, 2544-2546 (1969).
- 55) Lemieux, R.U. and Driguez, H.: The chemical synthesis of 2-acetamido-2-deoxy-4-Q-( $\alpha$ -L-fucopyranosyl)-3-Q-( $\beta$ -D-galactopyranosyl)-D-glucose, The Lewis a Blood-group Antigenic Determinant, J. Am. Chem. Soc., 97, 4063-4068 (1975).
- 56) Lemieux, R.U. and Driguez, H.: The chemical synthesis of 2-Q-( $\alpha$ -L-fucopyranosyl)-3-Q-( $\beta$ -D-galactopyranosyl)-D-galactose, The terminal structure of the blood-group B antigenic determinant, J. Am. Chem. Soc., 97, 4069-4075 (1975).
- 57) Liptak, A.; Jodal, I and Nanashi, P.: Stereroselective Ring-cleavage of 3-Q-Benzyl- and 2,3-Di-Q-benzyl-4,6-Q-benzylidene hexopyranose Derivatives with The  $\text{LiAlH}_4$ -

- $\text{AlCl}_3$  Reagent, Carbohydr. Res., **44**, 1-11 (1975).
- 58) Ljevakovic, D.; Srdanka, T. and Tomasic, J.: Selective Pivaloylation of 2-Acetamido-2-deoxy Sugars, Carbohydr. Res., **182**, 197-205 (1988).
  - 59) Magnusson, G; Noori, G.; Dahmen, J.; Frejd, T. and Lave, T.:  $\text{BF}_3$ -etherate induced formation of 2,2,2-trichloroethyl glycopyranosides, Selective visualization of carbohydrate derivatives on TLC plates, Acta. Chem. Scand. B **35**, 213-216 (1981).
  - 60) Mark, H.: Fifty Years of Cellulose Research, Cellulose Chem. Technol., **14**, 569-581 (1980).
  - 61) Matteucci, M.D. and Caruthers, M.H.: The use of zinc bromide for removal of dimethoxytrityl ethers from deoxynucleosides, Tetrahedron Letters, **21**, 3243-3246 (1980).
  - 62) Michael, A.: On The Synthesis of Helicin and Phenolglucoside, Am. Chem. J., **1**, 305-312 (1879).
  - 63) Miller, G.L.: Dean, J. and Blum, R.: A study of methods for preparing oligosaccharides from cellulose, Arch. Biochem. Biophys., **91**, 21-26 (1960).
  - 64) Miyamoto, T: Cellulose as Materials of Functional Polymers, Sen-I Gakkaishi, **42**, 25-30 (1986).
  - 65) Montogomey, E.M. and Richtmer, N.K.: The preparation and rearrangement of phenyl-glycoside, J. Am. Chem. Soc., **64**, 690-694 (1942).
  - 66) Nakamura, M. and Oki, M.: Restricted Rotation Involving The Tetrahedral Carbon XXXIV, Stable Rotamers of 9-

- Arylfluorences Carrying Oxygen-substituents in The Ortho Position of The Aryl Group, Bull. Chem. Soc. Jpn., 53, 3248-3251 (1980).
- 67) Nakatsubo, F.; Takano, T.; Kawada, T.; Someya, H.; Harada, T.; Shiraki, H. and Murakami, K.: Synthetic Studies of Cellulose I, Synthetic design and selection of the protective groups, Mem. Coll. Agric. Kyoto Univ., No.127, 37-47 (1985).
- 68) Nishino, S. and Ishido, Y.: Regioselective (Phenylcarbamoyl)ation of Polyhydroxy Compounds by Phenyl Isocyanate-Zinc Naphthematate, Carbohydr. Res., 155, 161-174 (1986).
- 69) Ogawa, T.; Nakabayashi, S. and Sasajima, K.: Synthesis of 2,2,2-trichloroethyl 3,6-di-O-benzyl-2-deoxy-2-phthalimido- $\beta$ -D-glucopyranoside and its reaction with glycosyl halides, Carbohydr. Res., 96, 29-39 (1981).
- 70) Ohno, Y.: The synthesis of cellulose by the method of organic chemistry, Sen-I Gakkaishi, 33, 87-94 (1977).
- 71) Ohno, Y.: Recent Progress in the Chemical Synthesis of Polysaccharides -On the Polysaccharides having Glucose as a Repeating Unit-, Koubunshi, 26, 110-116 (1977).
- 72) Oikawa, Y.; Yoshikawa, T. and Yonemitsu, O.: Specific removal of O-methoxybenzyl protection by DDQ oxidation, Tetrahedron Letters, 23, 885-888 (1982).
- 73) Paulsen, H.: Advances in Selective Chemical Synthesis of Complex Oligosaccharides, Angew. Chem. Int. Ed. Engl., 21, 155-173 (1982).
- 74) Paulsen, H.: Synthesis of Complex Oligosaccharide Chains

- of Glycoproteins, Chem. Soc. Rev., 15-45 (1984).
- 75) Perrin, D.; Dempsey, B. and Serjeant, E.P.: "pKa Prediction for Organic Acids and Bases", Chapman and Hall, London, 1981, p 60.
  - 76) Pigman, W. and Horton, D.: "The Carbohydrates, chemistry and biochemistry" Academic Press, New York and London, 1970.
  - 77) Pojer, P.M. and Angyal, S.J.: Methylthiomethyl ethers, General synthesis and mild cleavage, Protection of hydroxyl groups, Tetrahedron Letters, No.35, 3067-3068 (1976).
  - 78) Pojer, P.M. and Angyal, S.J.: Methylthiomethyl ethers, Their use in the protection and methylation of hydroxyl groups, Aust. J. Chem., 31, 1031-1040 (1978).
  - 79) Price, C.C.: "Organic Reactions, Vol.2", John Wiley & Sons Inc., New York, 1944, P 1.
  - 80) Satomura, S.; Iwata, T. and Sakata, Y.: Synthesis of p-nitrophenyl 6<sup>5</sup>-O-benzyl- $\alpha$ -maltopentaoside, A substrate alpha amylases, Carbohydr. Res., 176, 107-115 (1988).
  - 81) Schlubach, H.M. and Luhrs, L.: Ein Wirkung von Chlorowasserstoff auf Glucose, Synthese eines Polyglucan, Ann., 547, 73-85 (1941).
  - 82) Schmidt, R.R. and Michel, J.: Synthesis of linear and branched cellotetraose, Angew. Chem. Int. Ed. Engl., 21, 72-73 (1982).
  - 83) Schmidt, R.R.: O-( $\alpha$ -D-Glucopyranosyl) trichloroacetimidate as a glucosyl donor, J. Carbohydr.

- Chem., 4, 141-169 (1985).
- 84) Schmit, R.R.: New Methods for the Synthesis of Glycosides and Oligosaccharides -Are There Alternative to the Koenigs-Knorr Method?-, Angew. Chem. Int. Ed. Engl., 25, 212-235 (1986).
- 85) Seib, P.A.: 1,6-Anhydro-4-Q-benzyl- $\beta$ -D-glucopyranose, Carbohydr. Res., 8, 101-109 (1968).
- 86) Shapiro, D.; Rabinsohn, Y; Acher, A.J. and Diver-Haber, A.: Studies in the Ganglioside Series IV, Preparation of 2,3-Di-Q-acetyl-1,6-anhydro- $\beta$ -D-glucopyranose and Its Utilization in the Synthesis of Oligosaccharides, J. Org. Chem., 35, 1464-1467 (1970).
- 87) Sinay, P.: Recent Advances in Glycosylation Reactions, Pure & Appl. Chem., 50, 1437-1452 (1978).
- 88) Sugihara, J.M.: "Adv. Carbohydr. Chem. Biochem. Vol.8", Academic Press Inc, New York, 1953, p 2.
- 89) Takano, T.; Nakatsubo, F. and Murakami, K.: Synthetic Studies of Cellulose II, Synthesis of 1,4-substituted 2,3,6-tri-Q-acyl- $\beta$ -D-glucopyranosides, Mokuzai Gakkaishi, 33, 667-675 (1987).
- 90) Takano, T.; Nakatsubo, F. and Murakami, K.: Synthetic Studies of Cellulose V, Effect of the 3-Q- and 6-Q-substituent groups of the aglycon on  $\beta$ -glycosylation, Cellulose Chem. Technol., 22, 135-145 (1988).
- 91) Takano, T.; Harada, Y.; Nakatsubo, F. and Murakami, K.: Synthetic Studies of Cellulose VI, Effect of the substituent groups of the glycon on  $\beta$ -glycosylation, Cellulose Chem. Technol., submitted for publication.

- 92) Takano, T.; Harada, Y.; Nakatsubo, F. and Murakami, K.:  
Synthetic Studies of Cellulose VIII, Effect of the 3-O  
substituent group of the aglycon on  $\beta$ -glycosylation  
(2), Mokuzai Gakkaishi, submitted for publication.
- 93) Takano, T.; Nakatsubo, F. and Murakami, K.: A facile  $\beta$ -  
allylglycosylation in the presence of benzyl protective  
group using boron trifluoride etherate, Carbohydr. Res.,  
submitted for publication.
- 94) Takemoto, K.: Trial to synthesize cellulosic  
macromolecules, Sen-I Gakkaishi, 32, 101-106 (1976).
- 95) Takeo, K.; Yasato, T. and Kuge, T.: Synthesis of  $\alpha$ - and  
 $\beta$ -cellotriase hendecaacetates and of several 6,6',6"-  
tri-O-substituted derivatives of methyl  $\beta$ -cellotriase,  
Carbohydr. Res., 93, 148-156 (1981).
- 96) Takeo, K.; Okushino, K.; Fukuyama, K. and Kuge, T.:  
Synthesis of cellobiose, cellotriase, cellotetraose and  
lactose, Carbohydr. Res., 121, 163-173 (1983).
- 97) Takeo, K.; Nakaji, T. and Shinmitsu, K.: Synthesis of  
Lycotetraose, Carbohydr. Res., 133, 275-287 (1984).
- 98) Takeo, K. and Tei, S.: Synthesis of the Repeating Units  
of Schizophyllan, Carbohydr. Res., 145, 293-306 (1986).
- 99) Tomic, S.; Ljevakovic, D. and Tomasic, J.:  
Stereoselective hydrolysis of radiolabelled O-pivaloyl  
derivatives of methyl 2-acetamido-2-deoxy-D-  
glucopyranoside, Carbohydr. Res., 188, 222-227 (1989).
- 100) Tsuda, Y. and Yoshimoto, K.: Regioselective Syntheses  
of Acylated Mono-saccharides and Their Acyl Migration,

Yuki Gosei Kagaku, 42, 479-492 (1984).

- 101) Tsutsumi, H. and Ishido, Y.: Recent Advances in the Chemical Synthesis of Glycosyl Compounds, Yuki Gosei Kagaku, 38, 473-491 (1980).
- 102) Turner, S.: "The design of organic syntheses", Elsevier Scientific Publishing Company, Amsterdam-Oxford-New York, 1976.
- 103) Velluz, L.; Valls, J. and Mathieu, J.: Spatial arrangement and preparative organic synthesis., Angew. Chem. Int. Ed. Engl., 6, 778-789 (1967).
- 104) Wessel, H.P.; Inversen, T. and Bundle, D.R.: Acid-catalysed Benzylation and Allylation by Alkyl Trichloroacetimidates, J. Chem. Soc. Perkin Trans I, 2247-2250 (1985).
- 105) Whistler, R.L. and Wolform, M.L.: "Methods In Carbohydrate Chemistry Vol.II", Academic Press Inc, New York and London, 1963, p 211 and p 318.
- 106) Whistler, R.L. and Wolform, M.L.: " Methods In Carbohydrate Chemistry Vol.III", Academic Press Inc, New York and London, 1963, p 131.
- 107) Wulff, G and Rohle, G.: Results and Problems of Q-Glycoside Synthesis, Angew. Chem. Int. Ed. Engl., 13, 157-216 (1974).

## ACKNOWLEDGEMENTS

The author wishes to express his sincerest thanks to Professor Dr. Koji Murakami and Associate Professor Dr. Fumiaki Nakatsubo, Department of Wood Science and Technology, Faculty of Agriculture, Kyoto University, for their kindest guidance and encouragement during the entire course of this work.

The author is deeply grateful to Professor Dr. Takayoshi Higuchi, Research Section of Lignin Chemistry, Wood Research Institute, Kyoto University, and to Professor Dr. Tetsuo Koshijima, Research Section of Wood Chemistry, Wood Research Institute, Kyoto University, for their valuable suggestions and critical readings of the manuscript.

The author greatly appreciates to the staff of the Research Section of Lignin Chemistry, Wood Research Institute, Kyoto University, for the 200MHz  $^1\text{H}$ -NMR measurements, to Dr. Ritsuo Nishida, Pesticide Research Institute, Kyoto University, for the 90MHz  $^{13}\text{C}$ -NMR measurements, and to Associate Professor Dr. H. Imamura, Faculty of Agriculture, Kyoto University, for his useful discussion about  $\sigma$ -constant and PKa-value.

The author gratefully acknowledges the invaluable assistance of his co-researchers, Mr. Toshinari Kawada, Miss Yoshiko Harada, Mr. Toshiro Harada, Mr. Hiroshi Shiraki and Mr. Hiroyuki Someya.

Finally, Thanks are also due to Dr. Tatsuo Yamauchi, Dr. Makio Hasuike and other members of Laboratory of Forest



Products Chemistry, Department of Wood Science and Technology, Faculty of Agriculture, Kyoto University, for their kind help and discussion.